

INDEX, 1973, Nos. 1-4

NEW NAMES

Anthias boulengeri 200-201, 203, 206-207
Anthias conspicuus 200-209
Bertella idiomorpha 193-199
Boiga ocellata 580-586
Elaphe buisi 574-579
Eleutherodactylus elassodiscus 222-225
Gambusia amistadensis 210-221
Gasterosteus apodus 722-725
Lampetra minima 568-573

Liparis inquilinus 787-793
Melamphaes ebelingi 794-799
Parapristipomoides (new subgenus) 557
Pomacentrus exilis 565-568
Pristipomoides squamimaxillaris 557-559
Rhineuridae (new family) 705, 716-717
Rivulus stellifer 783-786
Spathorhynchus fossorium 704-720
Triathalassothia gloverensis 560-564

NEW AND ESTABLISHED NAMES

Abastor erythrogrammus 574
Ablepharus boutonii africanus 345-346
Aboma lucretiae 817
Abramis brama 45
Abudefduf zonatus 148
Acanthopagrus berda 353
Acanthurus chirurgus 609-610
coeruleus 609-610
triostegus sandvicensis 145
Acestrorhynchus 766-767
Achirus lineatus 25
Acipenser 360-361, 378-379
Acris crepitans 329
Acrocheilus alutaceus 396-400
alutaceus × *Richardsonius balteatus* 395-400
Acropoma 149
Adioryx diadema 145
Agamodon 712
Albula 768
Alepisaurus 41, 43
ferox 41-44
Alligator mississippiensis 635
sinensis 630
Alosa fallax 769
Ambassis 353
safga 353
Ambloplites rupestris 428-430
rupestris × *Chaenobryttus gulosus* 428-430
rupestris × *Enneacanthus obesus* 428-430
rupestris × *Lepomis gulosus* 428-430
rupestris × *Lepomis macrochirus* 428-430
rupestris × *Micropterus salmoides* 428-430
rupestris × *Pomoxis nigromaculatus* 428-430
Amblyopsis 378
rosae 110
spelaea 110
Amblyrhynchus 702

Ambystoma 676, 681
maculatum 189
mexicanum 680
Ameiurus nebulosus 110
Amia calva 653-657
Amphibolurus 111
barbatus 110, 338-339
pictus 338-339
Amphichthys 560, 562-564
cryptocentrus 560, 562-563
hildebrandi 562-563
Amphiprion 565
clarkii 203
polymnus 203
spp. 148
Amphisbaena 719
Amphiuma 805
tridactylum 459
Anchoa starksi 152
Ancylopsetta quadrocellata 154-155
Anguilla rostrata 364
Anolis 177, 270, 550, 552, 702
aeneus 264-271
carolinensis 267, 465-471, 623-624, 663, 667
grahami 471
lineotopis 267, 270
nebulosus 270
sagrei 267, 470-471
Anoplogaster cornuta 602-603
Anoplopoma fimbria 815
Anthias albofasciatus 201, 203
altus 201, 203
aplodactylus 203
bifasciatus 203
bilineatus 203
bimaculatus 201, 203
boulengeri 200-201, 203, 206-207
cheiropsilos 203
chrysostictus 203
cichlops 201, 203
clarkii 203
conspicuus 200-209
cooperi 202-203

- evansi* 200, 203, 207
formosus 203, 206
gibbosus 203
grunniens 203
heraldi 200, 203
huchtii 201, 203
hypselosoma 201, 203
japonicus 203
johnii 203
lepidolepis 203
maculatus 203
manadensis 201, 203
megalepis 203
mooreanus 202-203
mortoni 203
ornatus 203
pleurotaenia 202-203
rhodopeplus 203
squamipinnis 201, 203, 205
taeniatus 201, 203
townsendi 201, 203
vosmeri 203
- Anthiinae (new genus and species) 145**
Apeltes 379, 721, 725-734
 quadracus 722, 727, 730
Aphanius 75
Aphredoderus
 sayanus 377-379
Apogon
 maculiferus 145
 new sp. 145
 quadrisquamatus 89
 snyderi 145
Apsilus
 dentatus 149
Arapaima
 gigas 595
Archaeophelsuma 632
Archoplites
 interruptus 478
Argyrops
 spinifer 353
Arizona
 elegans 574
Ascapus
 truei 173-175, 759
Asterotheca 815-817
 alascana 816
 infraspinata 816
 pentacantha 816-817
Astyanax
 fasciatus 220, 232, 685
 mexicanus 348
Atelopus
 varius 190
Atropus 353
 atropus 353
Austrofundulus 783, 786
 dolichopterus 783, 786
 transilis 783
- Bagre* 378
Basiliscus 702
BathYGONUS 816
 alascanus 817
 infraspinatus 817
 nigripinnis 815-817
 pentacanthus 815-817
Bathyleptus 767
- Bathymyzon*
 bairdii 136
Batrachoides
 diemensis 562
 gilberti 560, 564
Batrachomoeus 562
 grunniens 560
Batrachus
 argentinus 563-564
 cryptocentrus 562
 grunniens 562
 rubigenes 562
Batrachthys 560, 562-563
 ophiocephalus 560, 563
Bellator
 brachychir 318
 egretta 318, 320
 militaris 318-319
 ribeiroi 318-319
 spp. 315-316, 318, 320-321
Belone 594
Belonesox
 belizanus 234
Bertella
 idiomorpha 193-199
Betta
 splendens 83
Bipes 705, 712, 715
Boiga
 cyanea 585-586
 cynodon 580-581, 586
 dendrophilus(a) 574, 585-586
 multimaculata 585-586
 ocellata 580-586
 ochracea 585-586
 quincunciata 585-586
 trigonata 585-586
 trigonatum melanocephala 801-802
Bothus
 spp. 156
Boulengerella 770
Brachydanio
 verio 673-681, 823-824
Brachylophus 702
Brevoortia 769
Brosomphyciops 32
Brosomphycis
 marginata 39
Brycon 763, 768-769, 773
Bufo 626
 americanus 343-344, 627
 arenarium 455
 boreas 342-344, 494, 496
 boreas halophilus 342-344
 bufo 495-496, 647, 650
 cognatus 103-107, 109-111, 322-329, 453-454, 458
 compactilis 322, 329
 debilis 442
 fowleri 440, 442-444
 marinus 343-344
 punctatus 458
 speciosus 322-329
 terrestris × *Bufo woodhousii fowleri* 61
 terrestris americanus × *Bufo woodhousii wood-*
 housii 61
 valliceps 189, 490-496, 627
 woodhousei fowleri 340-341, 490, 494-496
Bullisichthys
 caribbaeus 149-151

- Cadea* 712
Caesioperca
 thompsoni 145
Caiman
 crocodilus apaporiensis 630
 latirostris 630
 sclerops 804
Callisaurus 660, 701-702
Callochromis
 macrops 553
 macrops macrops 553
 macrops melanostigma 553
Campostoma 58
 anomalum 48, 54
 anomalum × *Notropis cornutus* 54
Canthigaster
 rivulatus 145
Carangoides 352-353
 armatus 352-354
 chrysophrys 353
 ciliarius 354
 cirrhosus 354
 citula 354
 ferdau 353
 malabaricus 353
 oblongus 353
 rupelli 354
 schlegeli 354
 uii 353
Caranx 352-353
 armatus 352-354
 ciliarius 354
 plumbeus 354
 schlegeli 354
Carassius 378
 auratus 46, 48, 110, 480
Carcharhinus
 albimarginatus 77
 falciformis 77
 galapagensis 76-77
 limbatus 77
 longimanus 77
 menisorrhah 76-83
Carcharodon
 carcharias 76
Careproctus
 melanurus 815
Carphophis
 vermis 472-476
Cataetx 39
 memorabilis 39
Catostomus
 occidentalis 478, 480-489
Centropyge
 argi 89
Centrotelma
 cryptomelan 515, 526-527
 geographica 527
Chaenobryttus
 gulosus 428-430
Chaetodon
 miliaris 145
 triangulum 146-148
 trifasciatus 146-148
 unimaculatus 148
Chanos 763
Characidium 766, 773-774
Chela 769
Chelonia
 mydas 305-311, 313-314
Chelydra
 serpentina 189, 505
Chirostoma 533-554
 aculeatum 534, 546, 548, 551
 arge 533-534, 545, 547-552
 attenuatum 547-548
 attenuatum attenuatum 549, 551
 attenuatum zirahuen 549-550
 bartoni 545, 548-549
 chapalae 546-547, 549, 551
 charari 545, 548-549
 compressum 547, 550
 consocium consocium 534, 546-547, 549, 551
 consocium reseratum 534, 546, 548-551
 estor 547-548
 estor copandara 546, 549-550
 estor estor 546, 549, 551
 grandocule 547, 549-551
 humboldtianum 533, 544, 546-547, 549-551
 jordani 533-534, 544-552
 labarcae 534, 546, 548-549, 551
 lucius 547, 551
 melanococcus 545, 547-549
 patzcuaro 546, 549-550
 promelas 547, 549, 551
 riojai 545, 547, 549
 sphyraena 547, 549, 551
Chologaster
 agassizi 110
Chorinemus 353
Chromis 565
 leucurus 145
Chrosomus
 eos 54
 eos × *Notropis cornutus* 54
 erythrogaster 54-59
 erythrogaster × *Notropis cornutus* 54-59
 oreas 54
 oreas × *Notropis cornutus* 54
Chrysemys
 picta 504, 513
Cichlasoma
 cyanoguttatum 220, 232
Citharichthys
 stigmaeus 155
Citula
 amphacantha 354
 armata 352, 354
 ciliaria 354
 plumbeus 353
Clemmys 360
 marmorata 21
Clinocottus 378-379
Clinostomus 51
 elongatus 45, 48, 50-51, 54
 elongatus × *Notropis cornutus* 54
 funduloides 54
 funduloides × *Notropis cornutus* 54
Clupanodon
 motius 736
Clupea
 finta 769
 melastoma 736-737
Cnemidophorus 111
 tigris 110, 446-451, 660
 velox 285
Coleonyx
 reticulatus 184
 variegatus 615-617

- Colostethus*
talamancae 278
Coluber
constrictor 476, 505-506, 508-509, 512-514, 574
Conolophus 702
Cophomantis
punctillata 515, 526-528
Cophosaurus 660
Coregonus
albula 186
clupeoides 186
hiemalis 186
maracana 186
sardinella 186
schinzi helveticus 186
Coryphaenoides
acrolepis 815
Corythophanes 702
Cottus
asper 480
gulosus 480, 487
Cotylin
nigripinnis nigripinnis 788
Couesius 51
plumbeus 48, 50, 369-370, 372
Crenichthys 378-379
Crocodylurus 740
Crocodylus
cataphractus 630
johnsoni 630
moreletii 630
siamensis 630
Crotalus
atrox 26-31
cerastes 26, 31
viridis 30, 285
cf. viridis 574
viridis lutosus 26
Crotaphytus 701-702
collaris 658, 660
wislizenii 658-661
Cryptobranchus
alleganiensis 331-337, 805-807
alleganiensis alleganiensis 331-337
alleganiensis bishopi 331-337, 805-807
Crythosaurus
mongoliensis 715
Ctenolucius 770
Ctenosaura 702
pectinata 267, 270, 431-434
Cubiceps 169
Culaea 589, 721, 724-734
inconstans 727, 730, 732
Curimata 766, 770
Cycleptus
elongatus 220
Cyclogaster
liparis 789
Cyclothone
acclinidens 600-601
signata 601
Cyclura 702
Cymatogaster 38
Cynolebias 783
spp. 786
Cynopoecilus 783
Cyprinodon 378-379
atrurus 682, 688
atrurus \times *Cyprinodon elegans* 688
bifasciatus 682
diabolis 137, 610-613
elegans 682-691
elegans \times *Cyprinodon macularius* 688
elegans \times *Cyprinodon nevadensis* 688
elegans \times *Cyprinodon radiosus* 688
elegans \times *Cyprinodon variegatus* 682-691
macularius 70-72, 75, 688
macularius \times *Cyprinodon variegatus* 688
milleri 366-369
nevadensis 137, 688
nevadensis \times *Cyprinodon variegatus* 688
nevadensis mionectes 137-138, 140
nevadensis pectoralis 137-140
radiosus 688
rubrofluvialis 68-75
rubrofluvialis \times *Cyprinodon variegatus* 688
salinus \times *Cyprinodon variegatus* 688
spp. 75
variegatus 682-691
Cyprinus 378, 764, 772
carpio 45, 48, 348, 480-481, 685, 824
Dactylagnus 116
mundus 115-119
Dactylanthias
aplodactylus 203
Dactyloscopus 116
new sp. 115
sp. 118
Daector
dowi 356-357
quadrizonatus 355-357
reticulatus 355-357
schmitti 355-357
Dallia 587
pectoralis 587
Dascyllus 565
aruanus 146, 148
Dasyatis
brevicaudatus 595
Decapturus 353
Dendrobates
pumilio 277-283
Desmognathus
fuscus 807
monticola 807
quadramaculatus 807
spp. 646
Diadophis
punctatus 476
Diaphus
taaningi 362-363
Dinematichthys 32, 39
ilucoeteoides 32-40
Diodon
holacanthus 609
hystrix 145
Diogenichthys 39
Dionda
episcopa 685
nubila 48
Diplodus 353
Diplomystes 762-763, 772, 775
Dipsas
cynodon 580
Dipsosaurus 702
dorsalis 100, 441, 504
Dolopichthys 193
pallatus 194-195, 198

Dormi
 latif
 Dracae
 Dryma
 cora
 Dryma
 mar
 Dryop
 nasu

Echis
 carin
 Egerni
 cum
 Elacat
 Elaph

buis
 diom
 gutt
 kans
 long
 nebr
 obso

57
 plio
 qua
 qua
 scal
 situ
 sub
 taen
 tria
 vulp

Elaps
 affi
 agla
 alie
 apia
 cora
 dias
 epis
 fulv
 fulv
 gua

Electr
 Eleotr
 gob
 pict
 pisc
 Eleuth
 bin
 bip
 crus
 disc
 elas

gras
 gue
 nas
 nigr
 octo
 par
 uni

Elops
 Emyd
 ma
 Emys
 orb
 Engro
 Engyo

- Dormitator*
latifrons 461-464
Dracaena 740
Drymarchon
corais 574
Drymobius
margaritiferus 574
Dryophis
nasutus 574
- Echis*
carinatus 801-802
Egernia
cunninghami 338-339
Elacatinus 609
Elaphe 574-575, 577, 634
buisi 574-579
dione 574
guttata 574-575, 577-579
kansensis 574-575, 578-580
longissima 574
nebraskensis 575, 578-580
obsoleta 505-506, 508-510, 512-514, 574-575, 577-579
pliocenica 575, 578-580
quadrivittata 505
quadrivittata 505
quadrivittata 505
scalaris 574
situla 574
subocularis 574-575, 577-579
taeniurus 574
triaspis 574-575, 577-579
vulpina 505-514, 574-575, 577-579
- Elaps*
affinis 1-2
aglaeope 2
alienus 2
apiatus 1-2
corallinus var. *crebrispunctatus* 1-2
diastema 1-2
epistema 2
fulvius 1
fulvius var. *sapperi* 2
guatemalensis 2
Electrophorus 763-764, 766, 770-771, 773
Eleotris
gobioides 169-170
picta 141
pisonis 141
Eleutherodactylus
binotatus 222, 224
biporcatus 222
cruralis 224
discoidealis 224
elassodiscus 222-225
granulosus 224
guentheri 224
nasutus 224
nigrovittatus 222, 224-225
octavioi 224
parvus 222
unistrigatus 222
Elops 768-769, 771, 773
Emydura
macquarrii 338-339
Emys
orbicularis 504, 513
Engraulis 39
Engyophrys
ciliaris 357-358
senta 357-358
Enneacanthus
obesus 428-430
Entonanthias 208
pascualis 207-208
Enyaliosaurus 702
Enyaloides 702
Eopelobates 190
Epinephelus 150
(Cephalopholis) fulvus 149
striatus 609
Eptatretus
deani 815
Erotelis
armiger 141
Eslopsarum
arge 550
jordani 550
Esox 378-379
lucius 124-132
Etelis 559
Etheostoma 377-379
(Etheostoma) blennioides 163
blennioides 162-164
blennioides pholidotum 163-164
(Oligocephalus) caeruleum 163
caeruleum caeruleum 163-165
(Nothonotus) camurus 163-165
(Catonotus) flabellare 163
flabellare flabellare 163-165
(Boleosoma) nigrum 163
nigrum nigrum 163-164
Eucyclogobius
newberryi 478
Eunneces 178, 703
egregius 345-347, 758
egregius similis 346-347
fasciatus 178-180
inexpectatus 178, 180
skiltonianus 285
Eumoea
atrocostatum 345
Eupomacentrus
fuscus 609
partitus 590-593
pictus 590-593
Eurycea
bilineata 807
bilineata wilderae 807
longicauda 340-341
longicauda melanopleura 341
lucifuga 340-341
Exoglossum 58
- Farancia*
abacura 574
Ficimia
olivacea 574
Floridichthys 75
Fundulus 35, 378-379
chrysotus 90-92, 808-809
confluentus 808-809
grandis 90-92
heteroclitus 369-370, 373
majalis 808
notatus 782
olivaceus 162, 782
zebrinus 71, 74

- Galaxias*
bongbong 363-365
Galeichthys
felis 153
Gambusia
affinis 19, 140, 217-220, 231, 250-252, 254-257,
 259-262, 417, 419-423, 478, 480-484, 486, 685
affinis affinis 138, 140
alvarezi 210-213, 215-216, 218-220
amistadensis 210-221, 252, 257-258, 260-261
atrora 253, 258-259
aurata 231, 252-253, 258-261
caymanensis 253, 258
gaigei 210-213, 215-220, 251-253, 258-262
geiseri 212, 219-220, 251-253, 255, 257-262, 685
georgei 219, 250-252, 257-262
heterochir 218, 220, 250-252, 254, 256, 258-262
hurtadoi 210-213, 215-216, 218-219, 252-253,
 257-260
krunholzi 218, 252-254, 258-260
longispinis 220
marshi 251, 253, 258-260
nicaraguensis 253, 258
nobilis 210-212, 215, 217-220, 251, 258, 260, 685
panuco 231, 253, 258-259
regani 231, 251, 253, 258-261
senilis 210, 212, 215, 217-218, 220, 253, 259
vittata 231, 253, 259
Gammaraia 249
Gasterosteus
aculeatus 417, 419-423, 427, 480, 487, 588-590,
 725-726, 728-731, 733-734
aculeatus aculeatus 722-723, 728, 731
aculeatus microcephalus 722-723, 731
aculeatus williamsoni 589, 722-723, 726, 728-
 729, 731-732
apodus 722-725, 734
doryssus 589-590, 722-723, 725
wheatlandi 722, 725-726, 728-729, 731-733
williamsoni 723
williamsoni leptosomus 723-724
Gaterin 353
Gehyra
variegata 617
Gephyrochromis 188
Gerrus
oyena 353
Gerrhonotus 703
coeruleus 699-700
kingi 699
leiocephalus 699
multicarinatus 699
sp. 698-700
Gila
bicolor 569
crassicauda 478
orcutti 55, 58, 417-427
orcutti × *Hesperoleucus symmetricus* 55, 58,
 417-427
Gillellus 116
Gilmorea
attenuatus 714
Ginglymostoma
cirratum 77, 606-608
Gladiunculus 733
Glyptocephalus
zachirus 815
Gnathanodon 353
speciosus 353
Gnathonemus
niger 595
petersi(i) 134-135, 595
tamandua 595
Gobiesox 378-379
Gobiomorphus
gobioides 169-170
huttoni 169-170
Gobiosoma
evelynae 609
genie 609
illecebrosum 609-610
(Elacatinus) illecebrosum 609
oceanops 609
prochilos 609
(Elacatinus) prochilos 610
Gobius
lucetiae 817-819
Goodea 549
Gorgasia
silneri 89
Gymnarchus
niloticus 595
Gymnothorax
sp. 145
Gymnotus 763-764, 766, 770-771, 773
sp. 134
Haemulon
sciurus 206
Halophryne 560, 562-563
diemensis 560
Haplochromis 187-188, 552-554
Hemidactylus
brookii 339-340
garnotii 339-340
Hemiergis 346
Hepsetus 767, 773
Hesperoleucus
symmetricus 55, 58, 417, 478, 480-489
symmetricus subditus 417-427
Hesperotestudo 360
Heterandria
bimaculata 234
Heterodon
plionasicus 574
Heterodontus
bonae-spei 165-167
francisci 166
galeatus 166
japonicus 165-166
philippi 165
portusjacksoni 165
quoyi 166
ramalheira 165-166
zebra 166
Heteropneustes
fossilis 370, 372
Heterotis
niloticus 595
Hiodon
alosoides 595
Holanthias
borbonius 203
chrysostictus 203
Holbrookia 660, 701-702
Holocentrus 353
Homesthes
caulopus 609

Hoplias
malin
Hoplo
Hoplo
Hoplo
litor
Hybog
hani
nuci
Hybof
aesti
stori
x-pu
Hydro
Hyla
albo
app
arbo
aviv
aviv
aviv
aviv
boan
calca
chry
chry
chry
chry
chry
cine
cine
cine
cruc
crypt
fabe
fasci
fem
fem
geog
geog
geog
geog
grat
lanc
lept
max
mul
para
pun
pun
rani
spec
squi
vers
Hytell
pun
Hypac
Hypoc
unif
Hypoc
torq
Hypso
gut
Hypsy
rubi
Hyster
trasi

Ichthy
Ichthy

- Hoplais*
malabaricus 785
Hoplocercus 702
Hoplosternum
litorale 785
Hybognathus
hankinsoni 47-48
nuchalis 48, 685
Hybopsis 51, 542
aestivalis 48, 220
storeriana 48
x-punctata 48
Hydrocyon 770
Hyla 681
albopunctata 528
appendiculata 515, 526, 528
arborea 673, 680
avivoca 411-415
avivoca × *Hyla chrysoscelis* 412-415
avivoca × *Hyla cinerea* 411-415
boans 515-516, 526, 529
calcarata 515-519, 522-525, 529, 531
chrysoscelis 411-414
chrysoscelis × *Hyla cinerea* 412, 414-415
chrysoscelis × *Hyla femoralis* 412, 414-416
chrysoscelis × *Hyla gratiosa* 412-415
chrysoscelis × *Hyla squirella* 412-416
cinerea 411-416
cinerea × *Hyla femoralis* 412-415
cinerea × *Hyla squirella* 412, 414-415
crucifer 61
cryptomelan 528
jaber 527-528
fasciata 515-519, 522-527, 529, 531
femoralis 411-415
femoralis × *Hyla squirella* 412-416
geographica 515-532
geographica geographica 528
geographica punctatissima 528
geographica var. semilineata 515, 526-528
gratiosa 411-416
lanciformis 525
leptoscelis 515, 522-523
maxima 527-528
multifasciata 525
pardalis 529
punctatissima 515, 527-528
punctillata 528
raniceps 525
spectrum 528-529
squirella 411-414
versicolor 61
Hylella
punctatissima 526-528
Hypacanthus 353
Hyporhamphus
unifasciatus 594
Hyporhina 714-716
Hypsigena
torquata 574
Hypsopsetta
guttulata 22-25
Hypsypops
rubicunda 145, 148
Hysterocarpus
traskii 478
Ichthyoborus 770
Ichthyokentema 769
Ichthyomyzon
castaneus 135
Ictalurus 378
catus 480-481
melas 220
nebulosus 480-481, 680
punctatus 220, 685
Icticus
ischanus 167
Idus 378
Iguana 111, 702
iguana 110, 296-304, 620-623
Ilisha
elongata 738
filigera 738
indica 736-738
macrophthalma 736
megaloptera 735-739
melastoma 735-739
motius 736
Iloglossus
calliurus 84, 90
helenae 84, 90
Iracundus
signifer 145
Jepsibaena
minor 714, 716-718
Jordanella 75
Kapparaia 249
Kyphosus 353
Labrisomus
kalisherai 597-598
Lacerta
teguixin 740-742
tupinambis 741-742
vivipara 189, 433
Laemactus 702
Lampanyctus 39
Lampetra
aepyptera 135
lethophaga 568-573
(Entosphenus) lethophaga 568-569
minima 568-573
(Entosphenus) minima 568-573
richardsoni 135
tridentata 568-573
(Entosphenus) tridentata(us) 136, 568-569
Lampropeltis 575
calligaster 574
getulus 498-503, 574
getulus brooksi 503
getulus floridana 501-503
getulus: floridana × *getulus* 501-503
getulus getulus 498, 500-503
getulus: getulus × *sticticeps* 498, 501
getulus sticticeps 497-503
pyromelana 574
triangulum 574
zonata 574
Latimeria
chalumnae 156-158, 813-814
Lavinia
exilicauda 478, 480-489

- Leiocephalus* 698, 700, 702-703
Leiocephalus 702-703
 carinatus 702
Leiopisma
 guichenoti 338-339
Leiopelma 173
 hochstetteri 173-175
Lepidochelys
 olivacea 306, 311-314
Lepidomeda
 albivallis 776-782
 altivelis 776-782
 mollispinis 776-782
 mollispinis pratensis 406-409
 vittata 776-782
Lepidosiren
 paradoxa 653
Lepidotrigla 316
 aspera 320
Lepisosteus 123, 653
 oculatus 119-123, 656-657
 osseus 656-657
 platyrhincus 656
 spatula 657
Lepomis
 auritus 131
 cyanellus 19, 220, 369-373, 417, 419-423, 425, 427, 478, 480-484, 486-488, 685
 gibbosus 110
 gulosus 428-430
 humilis 685
 macrochirus 124-132, 364, 428-430, 478, 480-484, 486, 685
 megalotis 363, 365
 microlophus 480-481
Leposternon 715, 719
Leptolebias 783
Lestophis 713-714
Lethrinus 353
 mahsena 353
Leuciscus
 baltatus 45
 leuciscus 45
Limanda
 ferruginea 155
Liolaemus 702, 755
Liparis
 atlanticus 787-789, 792
 cyclostigma 792
 inquinus 787-793
 koefoedi 792
 lineatus(a) 788
 liparis 787-792
 montagui 788
 tunicatus 792
 vulgaris 788, 791-792
Lobotes
 surinamensis 151
Lucasius
 damaeus 338-339
Lucioperca 186
Lutianus
 vaigensis 145
Lutjanus
 buccanella 149
 gibbus 200, 203
 johnii 203
 vaigensis 145
Lycodes
 diapterus 815
 Macolor 353
 Macrorhineura 714, 716-718
 Malacanthus
 plumieri 84, 89
 Marcgravia 562
 Marcgraviichthys 562
 Masticophis
 bilineatus 574
 flagellum 574, 803-804
 lateralis 575
 mentovarius 575
 taeniatus 285, 575
 Meda
 fulgida 776-782
 Medialuna
 californiensis 151
 Megalaspis 353
 Megalops
 atlanticus 140-141
 Megaprotodon
 strigangulus 146-148
 Melamphaes
 ebelingi 794-799
 lugubris 795
 microps 794-795, 798-799
 typhlops 798
 Melaniris 547
 crystallina 547
 Melanosuchus
 niger 630
 Menidia 378, 544, 547
 beryllina 544
 Merriamella
 doryssa 723-724
 Microlophichthys 198
 Micropterus
 dolomieu 480
 salmoides 220, 419, 425, 427-428, 478, 480-484, 486, 488
 Microstomus
 pacificus 815
 Micrurus
 affinis 1
 affinis affinis 1, 14
 affinis aglaeope 14
 affinis hippocrepis 14
 affinis mayensis 1-2, 14
 affinis stantoni 2
 alienus alienus 1, 14
 browni 2
 diastema 1-16
 diastema affinis 1
 diastema aglaeope 1
 diastema alienus 1
 diastema apiatus 1, 14
 diastema diastema 1
 diastema macdougalli 1-2, 14
 diastema sappieri 1
 elegans 2
 fulvius 2
 fulvius maculatus 2
 hippocrepis 1-2
 limbatus 2
 nirocinclus 2
 Mirolabrichthys 208
 dispar 208
 evansi 200, 203, 205, 207
 tuka 207-208
 tuka pascatus 205, 207-208
 tuka tuka 205, 208

- Moapa*
coriacea 408-409
Mollienesis
sphenops 229
Monitor
 (Tutor) *americanus* 741
meriani 741
Monodactylus 353
Monopeltis 715, 719-720
capensis capensis 718
Monotaxis 353
Morunasaurus 702
Moxostoma
aureolum 694
Mugil
cephalus 151
Mustelus
californicus 575-597
canis 76
Mylocheilus
caurinum 399
caurinum × *Ptychocheilus oregonensis* 399
Mylopharodon
conocephalus 478, 480-489
Myripristis 353
chryseres 145
Myxodagnus 116
- Naja* 634
Naso 353
Natrix
erythrogaster transversa 176-177
sipidon engelsi 498
sipidon fasciata 505
Necturus 805
Negaprion
brevirostris 77, 606
Neoceratodus
forsteri 504, 656-657
Neofundulus 783
Neoliparis
atlanticus 788
Neophelsuma 632
Neoseps
reynoldsi 758
- Oligopus* 33
ater 39
longhursti 33
Omosudis 41-43
lowei 41-44
Oncorhynchus
gorbuscha 58
gorbuscha × *Oncorhynchus keta* 58
keta 58
nerka 110-112, 130
Oneirodes 198
Ophisaurus 699
ventralis 500
Ophryoessoides 702
Opistognathus
aurifrons 84-90
Opsanus 562
astrifer 560, 564
beta 560, 564
hildebrandi 562
Opsariichthys 762, 769
- Opsopoeodus*
emiliae 47-48, 51, 161-163
Orestias 552, 554
Orthodon 358
microlepidotus 358-361, 478
Oryzias
latipes 369, 809-810
Osteocephalus
leprieurii 525
Osteoglossum
bicirrhosum 594-595
Ototriton 716-718
minor 716
solidus 714, 716
Oxyrrhites
typus 145
- Pachydactylus*
hasselquistii 634
Pantodon 594-595
buchholzi 595
Parabrotula 33, 38
Paragerrhonotus 699
Paralichthys
californicus 22-25
dentatus 155-156
Parapristipomoides 557-559
Parophrys
vetulus 155
Parrella 817-818
fusca 818-819
ginsburgi 818-819
lucretiae 818-819
macropteryx 818-819
maxillaris 818-819
spilopteryx 818
Pellona
ditchela 736
ditchoa 737
dussumieri 736-737
melastoma 736
motius 736
russelli 736
Pelomedusa 627-628
plioaenica 627
progaleata 627
subrufa 627
Pelusios 628
Pephrilus 169
simillimus 169
snyderi 169
Perca 378-379
flavescens 110, 124-132
Petalichthys
capensis 594
Petromyzon
marinus 136-137
Petrosaurus 701-702
Phelsuma 631-633
abbotti 632
abbotti parkeri 632
abbotti v-nigra 632
cepediana 632
flavicularis serraticauda 632
longinsulae pulchra 632
madagascariensis 632
madagascariensis grandis 632
mutabilis breviceps 632
newtonii 632

- quadriocellata bimaculata* 632
sundbergi 632
Phenacobius
mirabilis 48
Phillynodon
hubbsi 170
Phoxinus
eos 47-48, 51
erythrogaster 47-48, 51
neogaeus 47-48, 50-51
phoxinus 45, 348
Phrynosoma 360, 701-703
cornutum 701-702
coronatum 701
douglassi 698, 700-702
douglassi 701-702
m'calli 440-441, 701
modestum 701
platyrhinos 701
solare 701
Phyllobates
trinitatis 343
Pimephales 162
notatus 48
promelas 48, 685, 824
vigilax 48, 685
Pituophis 500, 575
catenifer 110
melanoleucus 575
Plagiotremus
spp. 153
Plagopterus
argentissimus 776-782
Platichthys
stellatus 155
Platyaster
indicus 737
macrophthalma 736
megalopterus 736-737
verticalis 737
Platymotus 702
Plectorhynchus 353
Plectranthias
anthioides 203
Plethodon
cinereus 441
dorsalis 807
Plica 702
Podocnemis
unifilis 181-183
Poecilia 262
butleri 226
formosa 226, 231, 350-352, 810-813
formosa × *Poecilia mexicana* 350-352, 813
limantouri 227
latipinna 140, 226, 231, 350
latipunctata 226, 231
mexicana 225-236, 350-352, 810-813
mexicana limantouri 225-236
mexicana mexicana 225-228, 232-236
mexicana: mexicana × *limantouri* 228, 231, 234-236
sphenops 226, 232-233
sulphuraria 226
Poecilopsis
gracilis 234
Pomacentrus 565, 567, 590
albolineatus 567
caeruleus 567
exilis 565-568
flavicauda 148
opisthostigma 567
partitus 89, 590-592
pictus 590-592
pulcherrimus 567
stigma 567
tablasensis 567
trichourus 567
variabilis 613-615
Pomadasy 353
grunniens 203
maculatus 203
Pomoxis
nigromaculatus 428-430, 824
Porichthys
bathoiketes 560, 564
Pranesus
insularum 151
Priacanthus 353
Prionobutis
koiomatodon 141-144
Prionotus
alatus 318
beanii 318
carolinus 317-320
evolans 318-320
griseus 318
ophrys 317-319
paralatus 318
pectoralis 317-319
punctatus 318
roseus 317-318
rubio 318
scitulus 316, 318
spp. 315-316, 318, 320-321
stearnsi 317-318
tribulus 317-318
Pristipomoides
freemani 559
squamimaxillaris 557-559
(Parapristipomoides) squamimaxillaris 559
Proctotretus 702
Protherapsis
trinitatis 277
Psenes 169
pellucidus 167-169
sio 167-169
Pseudacris
clarki 61, 495-496
nigrita 61
streckeri 61, 494-495
triseriata 61, 280, 490, 496
Pseudanthias
hypselosoma 203
Pseudemys
scripta 804
Pseudocerastes
persicus 801
Pseudopleuronectes
americanus 155-156
Pseudorhineura 714, 716
Pseudotropheus 188
Pterolebias
maculipinnis 783, 785-786
peruensis 783
zonatus 783, 786
Ptychocheilus 397, 764-766, 769
grandis 478, 480-489
oregonensis 399

- Ptyodactylus*
hasselquistii 634
Pungitius 589, 721, 724-734
haynesi 588-590
pungitius 722, 727, 729, 732
Pyloodictis
olivaris 220
Python
molurus 171-173
molurus bivittatus 171
molurus molurus 171
reticulatus 171-172

Rachovia 783
spp. 786
Raja
erinacea 238-249
lyllae 238, 249
ocellata 238-249
trachura 815
Rajella 238, 249
Rana 360, 681
aurora 18, 20-21
boas 527
boylli 18-21
cascaidae 415
catsebeiana 18-22, 103-107, 110-111, 277, 496
clamtans 277, 442, 496
esculenta 442, 458, 673, 680
pipiens 21, 103-107, 109-111, 174, 344, 441-443, 455, 626, 648
pipiens fisheri 21
pipiens pipiens 624-627
pretiosa 415
temporaria 348, 442, 496, 647-651
Rasbora
daniconius 348-349
Reinhardtius
hippoglossoides 156
Rhabdosargus
sarba 353
Rhineura 712-713, 715, 718-719
flavilata 759
floridana 704-705, 715-718, 758
hatcheri 718
Rhinichthys 162, 765
atratus 48, 50
cataractae 48, 50, 400-406
cataractae \times *Rhinichthys osculus* 395, 403-406
osculus 400-409, 419-423
osculus \times *Richardsonius balteatus* 395, 401-405
Rhinobatos
productus 76
Rhinoptera
bonasus 595
Rhyacotriton
olympicus 807
Richardsonius
balteatus 45, 396-400, 402-406
Rivulichthys 783
Rivulus 783
peruanus 783
stellifer 783-786
Rumula
spp. 153
Rypticus 150

Salminus 764

Salmo 35, 676, 680-681, 768, 771, 773
fario 110
gairdneri 420, 478, 480-483, 485-489
salar 348
salvelinus profundus 186
trutta 110, 480, 569
Salvador
merianae 741
Salvator
nigropunctatus 741
Salvelinus
fontinalis 569, 679
namaycush 110
profundus 186
willughbii 186
Samariscus
corallinus 145
Sardinops 39
caerulea 693
Sator 701-702
Sauromalus 110, 270, 702
hispidus 110
obesus 93-102, 440
Scaphiopus
bombifrons 60-67, 435, 452-459
bombifrons \times *Scaphiopus hammondii* 60-67
couchi(i) 435-444, 452-459, 627
hammondii(i) 60-67, 103-114, 435-444, 452-459, 627
hurteri 495
Scarus
coelestinus 609
croicensis 609
guacamaia 609
vetula 609
Sceloporus 698, 700-701, 703
Sceloporus 270, 276, 701-703
graciosus 284-295, 433
gramnicus 270
jarrovi 272-276, 450
cf. magister 701
malachiticus 276
occidentalis 341-342, 433, 662-671
olivaceus 759
spp. 660
undulatus 285, 342, 470, 748, 751, 756
undulatus consobrinus 756
undulatus undulatus 756
virgatus 660, 662, 756
woodi 746-760
Schultzea 149
Sciaena 353
armata 352-354
Sclerophages 595
formosus 595
leichardtii 595
Scolopsis 353
bilineatus 203
vosmeri 203
Scomber 353
lysan 353
Scomberesox
saurus 594
Scomberoides 353
lysan 151-154
Sebastodes 38
Sebastolobus
alascanus 815
altivelis 815
Selar 353

- Semotilus* 51, 58
atromaculatus 45, 48, 50-51, 54, 58
corporalis 54
margarita 48, 50-51
Seps
marmoratus 743
Seriola 353
Seriolina 353
Serranus 41, 149-150
cichlops 203
(Anthias) squamipinnis 203
tortugarum 89
Siganus 353
rostratus 604-606
spinus 604-606
Simpsonichthys 783
Siren 805
Smaris
mauri 317
vulgaris 317
Smilodonichthys 360
Sparisoma
rubripinne 609
viride 609-610
Spatorhynchus
jossorium 704-720
Sphenodon 111, 662
punctatum(s) 110, 504, 617-620
Sphenomorphus
taeniolatus 338-339
Sphyrna
lewini 595
sp. 77
tiburo 76
Spinachia 721, 723, 726-734
vulgaris 722, 727, 730-732
Stegostoma
fasciatum 595
Stenocercus 702
Sternotherus 476
odoratus 305-306, 309-312, 314
Stilosoma
extenuatum 758
Stizostedion 186, 378-379
Stolephorus
purpureus 152-153

Taenianotus
triacanthus 145
Taenioconger
hassi 89
Takydromus
tachydromoides 450
Tantilla
relicta 758
Taricha 360
Tejus
rufescens 743
Telescopus
fallax 801
rhinopoma 801
Terapon 353
Teratolepis
fasciatus 801
Terrapene
carolina 313
Testudo
ibera 513

Thalassophryne
amazonica 356
maculosa 355
quadrizonatus 355-356
Thamnophis 178, 476
radix 476
sirtalis 180
sp. 574
Thymallus
arcticus 819-822
Tilapia 188
Tiliqua 111
nigrolutea 338-339
rugosa 338-339
scincoides 110
Tomuropeltis 715, 719
Torpedo
californica 76
Trachinotus 353
Trachurus 353
Triakis
semifasciata 596
Triathalassothia 564
argentinus 560, 563-564
devincenzii 563
gloverensis 560-564
Tridentiger
trigonocephalus 143
Trigla 316
gurnardus 317
Trigonectes
strigabundis 783
Trogonophis 712
Tropidurus 702
albemarlensis 671
Tupinambis
duseni 740-741, 743
monitor 740-742
nigropunctatus 740-742
rufescens 740-745
teguixin 740-745
teguixin sebastiani 740, 743
Tylosurus
acus 594
Typhlichthys
subterraneus 110

Ulua
mandibularis 353
Uma 633, 701-702
inornata 100, 662
notata 100
scoparia 100
Umbra 587-588
krameri 587-588
limi 587-588
praekrameri 587
pygmaea 587-588
weileri 587
Uranoscodon 702
Urocentron 702
Urosaurus 701-702
ornatus 270
Uta 111, 270, 701-702, 755, 758
stansburiana 110, 285, 287, 292, 450, 470, 658, 660

- Varanus* 111
 - gouldii* 110
 - niloticus* 742
 - salvator* 742
- Verriculus*
 - sanguineus* 145
- Vinciguerria* 39
- Vipera*
 - berus* 476
 - persica* 801
- Vorhisia*
 - vulpes* 692-697
- Xantusia* 476
 - vigilis* 662
- Xeneretmus* 815-816
 - latifrons* 815
 - leiops* 814-815
- Xenochirus*
 - pentacanthus* 815
- Xenomystus*
 - nigri* 595
- Xiphophorus* 158-161, 235
 - couchianus* 232
 - helleri* 158-161, 234
 - helleri* × *Xiphophorus maculatus* 159-161
 - maculatus* 158-161, 234
 - montezumae* 232
 - variatus* 232-234
 - xiphidium* 232
- Zacco* 769
- Zygonectes* 782

SUBJECT INDEX

ACTIVITY (see BEHAVIOR)

AGE AND GROWTH (see GROWTH)

AGONISTIC DISPLAY (see BEHAVIOR)

ANATOMY

embryonic, *Dinematichthys ilucoeteoides* 33-38.
eye septum, *Osteoglossum bicirrhosum* 594-595.

functional (jaw), *Latimeria chalumnae* 156-158.

infraoral laminae, *Petromyzon marinus* 136-137.

reproductive system, *Omosudis lowei*, *Alepisaurus ferox* 41-44; *Bullisichthys caribbaeus* 149-151.

respiratory structures, Dactyloscopidae (*Dactylagnus*, *Gillellus*, *Dactyloscopus*, *Myxodagnus*) 115-119.

swimbladder and associated structures, Triglidae (*Bellator* [four species], *Prionotus* [14 species]) 315-321.

teeth, *Heterodontus bonae-spei* (= *Heterodontus portusjacksoni*) 165-167.

ANOMALIES, *Hypsopsetta guttulata*, *Paralichthys californicus* 22-25.

ASSOCIATIONS, Fish species associations in Sierra Nevada foothills, California 478-490; *Labrisomus kalisherae* (with sea anemone) 597-598; *Gobiosoma (illecebrosus, prochilos)* (feeding on ectoparasites of larger fishes) 609-610; *Rivulus stellifer* 786; *Liparis inquilinus* (with scallops) 787-794.

BATHYMETRIC DISTRIBUTION, *Cyclothone acclinidens* 600-601.

BEHAVIOR, *Sphenodon punctatus* (in T-maze); *Iguana iguana* (post-nesting movements from nesting aggregations) 620-622; *Fundulus (chrysotus, confluentus)* (response to drought) 808-809.

Agonistic displays and attacks on man, *Caracharhinus menisorrh* 76-83.

Breeding, *Chrosomus erythrogaster* and *Notropis cornutus* 58-59; *Cyprinodon rubrofluviatilis* 68-75; *Gambusia (heterochir, affinis, georgei, geiseri, regani, gaigei, marshi)* (anal spot variation and effects on courtship) 250-263.

Burrowing, *Opistognathus aurifrons* 84-90; *Scaphiopus hammondi* (during forced activity) 105-107.

Courtship, *Gambusia (heterochir, affinis, georgei, geiseri, regani, gaigei, marshi)* (anal spot variation and effects on courtship) 250-263.

Displays, *Anolis aeneus* (related to social organization) 264-271.

Feeding, *Cyprinodon rubrofluviatilis* 68-75; *Scomberoides lysan* and *Oligoplites* (2 spp.) (scale-feeding behavior in young) 151-154. *Ancylosetta quadrocclata*, *Pseudopleuronectes americanus*, *Paralichthys dentatus*, *Limanda ferruginea*, *Citharichthys stigmatheus*, *Parophrys vetulus*, *Platichthys stellatus* (fin use in relation to feeding) 154-156; *Bufo boreas* (social behavior during feeding in immature individuals) 342-344; *Anoplogaster cornuta* 602-603; *Ginglymostoma cirratum* (suction feeding) 606-608.

Forced activity, *Scaphiopus hammondi* (during burrowing) 105-107.

Nest construction, *Lampetra aepyptera* 135.

Orientation, *Crotalus atrox* (relation to) 26-31.

Oxygen, *Lepisosteus oculatus* (effect of dissolved oxygen on locomotor reactions) 119-123; *Esox lucius*, *Perca flavescens*, *Lepomis macrochirus* (response to low oxygen concentrations under winterkill conditions) 124-133.

Predation, *Eumeces fasciatus* (attacks in young triggered by chemical cues) 178-180.

Residence time, *Megaprotodon strigangulus*, *Chaetodon (trifasciatus, triangulum, immaculatus)*, *Pomacentrus flavicauda*, *Amphiprion* spp. (duration of residence on "home" reefs) 145-149.

Social, *Anolis aeneus* 264-271; *Bufo boreas* (social behavior during feeding in immature individuals) 342-344.

Territorial, *Sauromalus obesus* 93-102; *Dendrobates pumilio* 277-283.

Visual discrimination, *Podocnemis unifilis* 181-183.

Vocalizations, *Dendrobates pumilio* (effects on territorial behavior) 277-283.

BIOCHEMISTRY, *Xiphophorus (helleri, maculatus)* (isocitrate dehydrogenase) 158-161; *Natrix erythrogaster transversa* (serum proteins and nutritional state) 176-178; *Eumeces fasciatus* (chemical release of prey attack) 178-181; *Bufo (cognatus, speciosus)* (protein polymorphism, genic heterozygosity) 322-329.

BLOOD, *Scaphiopus hammondi* (lactate levels) 105-109; *Cryptobranchus alleganiensis (alleganiensis and bishopi)* (comparative hematology) 331-337; eleven lizard species (method of obtaining blood from eye) 338-339;

Scaphiopus (couchii, hammondii) 453;
Elaphe (obsoleta, vulpina), *Coluber constrictor* (serum immune responses) 504-514.
BREEDING (see REPRODUCTION)
BUOYANCY, *Dormitator latifrons* (mode of inflation and function of gas bladder) 461-464.
BURROWING (see BURROWS)
BURROWS (and BURROWING), *Opistognathus aurifrons* (construction) 86-88; *Opistognathus aurifrons*, *Malacanthus plumieri*, *Ioglossus (calliurus, helenae)*, *Nystactichthys halis* 84-90; *Scaphiopus hammondii* 105, 107-108, 112-113.
CALLS (see SOUND PRODUCTION)
CHROMOSOMES (see KARYOTYPES)
COLOR (see PIGMENTATION)
COURTSHIP (see BEHAVIOR)
DDT, *Rana temporaria* (response of tadpoles) 647-652.
DESCRIPTIONS (including **DIAGNOSES**) (not including new species), *Micrurus diastema* 2-3; *Heterodontus bonae-spei* (= *Heterodontus portusjacksoni*) (based on jaws of holotype) 165-166; *Anthias bouleengeri* 206-207; *Mirolabrichthys (evansi, tuka tuka, tuka pascualis, dispar)* 207-208; *Poecilia mexicana (mexicana and limantouri)* 227-229, 232-233; *Raja (erinacea, ocellata)* 240-247; *Daector quadrizonatus* 355-356; *Engyophrys senta* 358; *Oligocottus maculosus* (larvae) 374-376; *Acrocheilus alutaceus* × *Richardsonius balteatus*, *Rhinichthys osculus* × *Richardsonius balteatus*, *Rhinichthys osculus* × *Rhinichthys cataractae* 399-408; *Hyla* (hybrid tadpoles involving six species: *cinerea*, *avivoca*, *gratiosa*, *squirella*, *femoralis*, *chrysoscelis*) 412-416; *Lampropeltis getulus sticticeps* 500-502; *Hyla (calcarata, fasciata, geographica)* 521-530; *Eupomacentrus (partitus, pictus)* 591-593; *Mustelus californicus* (albinos) 595-596; *Vorhisia vulpes* (fossil otoliths) 692-697; *Gerrhonotus* sp., ?*Sceloporus*, *Phrynosoma* ?*douglasi*, ?*Leiocephalus* (fossils) 699-703; *Gasterosteus* (apodus, doryssus, aculeatus aculeatus, a. microcephalus, a. williamsoni) (both fossil and Recent forms) 722-734; *Ilisha (megalopectera, melastoma)* 736-739; *Tupinambis (teguixin, rufescens)* 740-743; *Sceloporus woodi* 750; *Bathyaenonius (nigripinnis, pentacantha)* 816; *Parrella (lucetiae, fusca, ginsburgi, macropteryx, maxillaris)* 818-819; *Thymallus arcticus* 820-822.

DIAGNOSES (see **DESCRIPTIONS**)
DISCRIMINATION, *Podocnemis unifilis* (visual) 181-183.
DISPERSAL, *Brotulidae* (mode of dispersal of pelagic larvae) 39; *Poecilia mexicana* (northeastern Mexico) 234-236; *Chiostoma* (18 species) (in central Mexico) 533-544; *Sceloporus woodi* (dispersal during Pleistocene) 746, 755-758.
DISTRIBUTION MAPS, *Micrurus diastema* 4, 15; *Cyprinodon nevadensis pectoralis* 138; *Bertella idiomorpha* 197; *Poecilia mexicana* (northeastern Mexico only) 228; *Orthodon microlepidotus* and fossil form) 359; *Lampropeltis getulus* (coastal North Carolina only) 498; *Hyla calcarata* 525; *Hyla fasciata* 527; *Hyla geographica* 530; genus *Chiostoma* (18 species) 535; *Chiostoma (jordani, arge, riojai, bartoni, charari, melanococcus)* 545; *Chiostoma (humboldtianum, aculeatum, labarcae, consocium consocium, consocium reseratum, estor estor, estor copandaro)* 546; *Tupinambis (teguixin and rufescens)* 744; *Sceloporus woodi* 747; *Lepidomeda (vittata, mollispinis, altivelis, albivialis)*, *Meda fulgida*, *Plagopterus argentsimus* 777.
DISTRIBUTIONS (see **GEOGRAPHIC RANGES**)
DNA, *Poecilia (formosa, mexicana)* 810-813.
DROUGHT, *Fundulus (chrysotus, confluentus)* (response to) 808-809.
ECOLOGY, *Micrurus diastema* 15-16; *Rana catesbeiana* [competition with native *Rana (aurora, boylei)*] 18-20; *Notropis cornutus*, *Chrosomus erythrogaster* and hybrids 58-59; *Opistognathus aurifrons* 84-90; *Sauromalus obesus* 93-102; *Lampetra aepyptera* (nest construction) 135; *Cyprinodon nevadensis pectoralis* 138-140; *Megalops atlanticus* (larvae) 140-141; *Prionobutis koolamotodon* (factors precipitating introduction to Panama) 141-144; Twenty species in sunken vessel (Hawaii) 144-145; *Megaprotodon strigangulus*, *Chaetodon (trifasciatus, triangulum, immaculatus)*, *Abudefduf zonatus*, *Amphiprion* spp., *Dascyllus aruanus*, *Hypsypops rubicunda*, *Pomacentrus flavicauda* (duration of residence on home reefs) 145-149; *Scomberoides lysan*, *Oligoplites (refulgens, saurus inornatus)* 151-154; *Gambusia (amistadensis, geiseri, nobilis, alvarezi, gaigei, hurtadoi)* 219-220; *Eleutherodactylus elassodiscus* 224; *Poecilia mexicana (mexicana and limantouri)* 230-234; *Sceloporus*

jarrovi (reproductive effort and winter activity) 272-276; *Dendrobates pumilio* (vocalizations and territorial behavior) 277-283; *Sceloporus graciosus* 284-295; *Gambusia* (*affinis*, *georgei*, *heterochir*) (courtship) 250-263; *Anolis aeneus* (displays and social organization) 264-271; *Bufo boreas* (social behavior in young) 342-344; *Rasbora daniconius* (seasonal changes in pituitary gland) 348-349; *Galaxias bongbong* 363-365; *Cyprinodon milleri* (thermal environment) 366-369; *Gila orcutti*, *Hesperoleucus symmetricus* and hybrids 419-423; *Scapthiopus* (*couchii* and *hammondi*) (beneath ground) 452-459; native and introduced fishes in central California (21 species total) 478-489; *Bufo valliceps* (olfactory orientation) 490-496; *Lampropeltis getulus sticticeps* (outer banks of North Carolina) 497-503; *Hyla calcarata* 523; *Hyla fasciata* 526; *Hyla geographica* 529-530; *Labrisomus kalisherae* (associated with sea anemone) 597-598; *Thorogobius ephippiatus* (habitats off southern England) 598-600; *Cyclothone acclinidens* (ecology and bathymetric distribution) 600-601; *Gobiosoma* (*illecebrosus*, *prochilos*) (parasite pickers) 609-610; *Cyprinodon diabolis* (reproductive cycle) 610-613; *Pomacentrus variabilis* (effects of hurricane) 613-614; *Iguana iguana* (post-nesting movements) 620-622; *Pelomedusa subrufa* (correlation with abundance in fossil record) 627-629; *Notophthalmus viridescens* (life history variation) 641-647; *Cyprinodon* (*elegans*, *variegatus*, and hybrids) 684-687; *Tupinambis* (*teguixin*, *rufescens*) 744; *Sceloporus woodi* 746-760; *Rivulus stellifer* 785-786; *Liparis inquilinus* 787-788.

EGGS, *Leiopelma hochstetteri* (nuclear constitution of oocytes) 173-175.

ELECTROCARDIOGRAPHS, Squamate reptiles (evidence of a contractile bulbus cordis) 802-805.

ELECTROPHORESIS, *Bufo* (*cognatus*, *speciosus*) (protein polymorphism, gene heterozygosity, and divergence) 322-329.

EMBRYOLOGY, *Dinematichthys ilucoeteoides* 32-40; *Python molurus* (effects of temperature on development) 171-173; *Leiopelma hochstetteri* (nuclear constitution of oocytes) 173-175; *Iguana iguana* (embryonic growth) 296-304; *Hemidactylus garnotii* (skin heterograft rejection) 339-340; *Oligocottus maculosus* (description of laboratory-reared larvae) 373-377; *Aphredoderus sayanus* (comments on yolk circulation patterns) 377-379;

Mustelus californicus (albinism in embryos) 595-597; *Cyprinodon diabolis* (egg development) 610-613; *Brachydanio rerio* (development of Mauthner neurons in embryos and larvae) 673-681; apparatus for development of aquatic embryos 809-810.

EVOLUTION (see **TAXONOMY** in part), North American Cyprinidae (trends in pharyngeal tooth evolution) 50-52; *Poecilia mexicana* (northeastern Mexico) 234-236; *Chiostoma* (18 species) (Mexico) 533-556; *Elaphe* (seven species) 579; *Umbra* spp. (North America) 587-588; *Gasterosteus* spp. 589-590; pelomedusid turtles (correlation between ecology and abundance in fossil record) 627-629; family Rhineuridae (and *Spathorhynchus fossorium*) 704-720; family Gasterosteidae (Pliocene fossils and Recent) 721-735; *Sceloporus woodi* 746-761; Cypriniform fishes (special reference to feeding structures) 761-776; Plagopterin cyprinids (chromosomes) 776-782.

EXOTICS, *Prionobutis koilomatodon* (in Panama) 141-144.

EXTINCTIONS, *Gambusia amistadensis* 210-221.

FEEDING (see **FOOD**)

FINS, flatfishes (seven species) (fin use in relation to feeding) 154-156.

FOOD (including **FEEDING**), *Cyprinodon rubrofluviatilis* (feeding behavior) 69-70; *Sauro-malus obesus* (diet) 93-102; *Scomberoides lysan*, *Oligoplites* (*refulgens*, *saurus inornatus*) (food and feeding behavior, particularly as relates to scales) 151-154; *Ancylor-setta quadrocellata*, *Limanda ferruginea*, *Paralichthys dentatus*, *Pseudopleuronectes americanus*, *Citharichthys stigmaeus*, *Parophrys vetulus*, *Platichthys stellatus* (food and feeding behavior) 154-156; *Gambusia amistadensis* (food) 220; *Poecilia mexicana limantouri* (food) 231; *Anolis aeneus* (feeding behavior) 266; *Bufo boreas* 342-344; *Ablepharus boutonii africanus* (food habits) 345-346; *Gila orcutti*, *Hesperoleucus symmetricus* (and hybrids), *Gasterosteus aculeatus*, *Rhinichthys osculus*, *Gambusia affinis*, *Micropterus salmoides*, *Lepomis cyanellus* 421-423, 426, 427; *Ctenosaura pectinata* 431-434; *Pomacentrus exilis* 568; *Anoplogaster cornuta* (feeding behavior) 602-603; *Siganus* (*rostratus*, *spinus*) (food preference of juveniles) 604-606; *Ginglymostoma cirratum* (suction feeding) 606-608.

FOSSILS, *Orthodon microlepidotus* (Pliocene and Pleistocene) 358-361; *Elaphe buisi* (Pliocene) 574-579; *Umbra limi* (Pleistocene) 587-588; *Gasterosteus aculeatus* (Pliocene) 588-590; *Pelomedusa* (correlation between ecology and abundance in Pleistocene fossil record) 627-629; *Vorhisia vulpes* (otoliths from upper Cretaceous) 692-697; *Gerrhonotus* sp., ?*Sceloporus*, *Phrynosoma* ?*douglassi*, ?*Leiocephalus* (Miocene) 698-703; *Spathorhynchus fossorium* (middle Eocene; discussion of fossil history of family Rhineuridae) 704-720; *Gasterosteus* (*apodus*, *doryssus*) (Pliocene) 721-734.

FUNCTION (also see **ANATOMY**), *Latimeria chalumnae* (jaws) 156-158.

GAS BLADDER, *Prionotus* (14 species), *Bellator* (four species) (morphology) 315-321; *Dormitator latifrons* (function) 461-464.

GAS EXCHANGE (see **PHYSIOLOGY**)

GENETICS (see **KARYOTYPES**)

GEOGRAPHIC LOCALITIES

Africa, *Pelomedusa* spp., *Pelusios* spp. 627-629.

Alaska, *Thymallus arcticus* 819-822.

Argentina (see South America)

Arizona, *Scaphiopus hammondi*, *Rana pipiens* 103-114; *Sceloporus jarrovi* 272-276; *Scaphiopus* (*couchii*, *hammondi*) 435-444; *Scaphiopus* (*bombifrons*, *hammondi*), *Bufo cognatus* 452-459; *Lepidomeda* (*mollispinis*, *vittata*), *Meda fulgida*, *Plagopterus argentinus* 776-782.

Atlantic Ocean (eastern), *Petromyzon marinus* (off Ireland) 136-137.

Atlantic Ocean (western), *Raja* (*erinacea*, *ocellata*) (various localities) 239-243; *Prionotus* (14 species), *Bellator* (four species) (various localities) 315-321; *Engyophrys senta* 357-358; *Diaphus taaningi* (Caribbean Sea) 362-363; *Liparis inquilinus* (Gulf of St. Lawrence to Cape Hatteras, North Carolina) 787-793; *Melamphaes ebelingi* (off Bermuda) 794-799.

Australia, *Megaprotodon strigangulus*, *Chaetodon* (*trifasciatus*, *triangulum*, *immaculatus*), *Pomacentrus flavicauda*, *Amphiprion* spp. 145-149; *Heterodontus portusjacksoni* 165-167; *Amphibolurus* (*barbatus*, *pictus*), *Tiliqua* (*rugosa*, *nigrolutea*), *Egernia cunninghami*, *Leiolopisma guichenoti*, *Sphenomorphus taeniolatus*, *Oedura marmorata*, *Lucasius damaeus*, *Emydura macquarrii* 338-339; *Galaxias bongobong* 363-365.

Bahamas, *Opistognathus aurifrons* 84-90; *Ginglymostoma cirratum* 606-608.

Barbados, *Eupomacentrus partitus* 590-593.

Belize (see British Honduras)

Bolivia (see South America)

Brazil (also see South America), *Eupomacentrus pictus* 590-593.

British Honduras (= Belize), *Micrurus diastema* 3; *Triathalassothia gloverensis* 560-564.

California, *Rana* (*catesbeiana*, *aurora*, *boylei*) 18-22; *Paralichthys californicus*, *Hypospsetta guttulata* 22-25; *Sauromalus obesus* 93-102; *Hypsypops rubicunda* 148; *Bufo boreas halophilus* 342-344; *Orthodon microlepidotus* 358-361; *Cyprinodon milleri* 366-369; *Oligocottus maculosus* 373-377; *Gila orcutti* × *Hesperoleucas symmetricus*, *Gambusia affinis*, *Gasterosteus aculeatus*, *Lepomis cyanellus*, *Micropterus salmoides* 417-427; 21 species (nine native, 12 introduced) 478-489; *Pungitius haynesi* (= *Gasterosteus aculeatus*) 588-590; *Mustelus californicus* 595-597; *Cyclothone acclinidens* 600-601; *Anoplogaster cornuta* 602-603; *Sceloporus occidentalis* 662-671.

Canal Zone, *Dormitator latifrons* 461-464; *Iguana iguana* 620-622.

Caribbean Sea, *Bullisichthys caribbaeus* (Sombrero Cay Bank) 149-151.

Caroline Islands, *Pomacentrus exilis* 565-568.

Cayman Islands, *Gambusia caymanensis* 250-263.

Colombia (also see South America), *Daector* (*dowi*, *quadrizonatus*) 355-357.

Comoro Islands, *Latimeria chalumnae* 813-814.

Costa Rica, *Dendrobates pumilio* 277-283; *Chelonia mydas* 305-314; *Daector schmitti* 356.

Easter Island, *Pristipomoides squamimaxillaris* 557-559.

Ecuador (also see South America), *Eleuthero-dactylus elassodiscus* 222-224.

England, *Thorogobius ephippiatus* 598-600; *Rana temporaria*, *Bufo bufo* 647-651.

Florida, *Megalops atlanticus* 140-141; *Eumeces egregius* 346-347; *Eupomacentrus partitus* 590-593; *Pomacentrus variabilis* 613-614; *Sceloporus woodi* 746-760; *Fundulus* (*confluentus*, *chrysotus*) 808-809.

French Guiana (see South America)

French Polynesia (specifically Rapa), *Pristipomoides squamimaxillaris* 557-559.

Georgia, *Ancyplossetta quadrocellata* 154-156.

Great Britain (see England)

- Guam, *Siganus (rostratus, spinus)* 604-606.
 Guatemala, *Micrurus diastema* 3.
 Guyana (see South America)
 Hawaii, *Omosudis lowei*, *Alepisaurus ferox* 41-44; Twenty species of reef fishes 144-145; *Scomberoides lysan* 151-154.
 Honduras, *Gambusia nicaraguensis* 250-263.
 Illinois, *Notropis cornutus*, *Chrosomus erythrogaster*, and hybrids 54-59; *Elaphe (obsoleta, vulpina)*, *Coluber constrictor* 504-514.
 India, *Rasbora daniconius* 348-349; *Ilisha (megaloptera, melastoma)* 735-739.
 Indiana, *Elaphe (obsoleta, vulpina)*, *Coluber constrictor* 504-514.
 Indian Ocean, *Anthias (boulengeri, conspicuus)* 200-207.
 Iran, *Boiga trigonatum melanocephala*, *Echis carinatus* 801-802.
 Ireland, *Petromyzon marinus* 136-137.
 Kenya, *Hemidactylus (brookii, garnotii)* 339; *Ablepharus boutonii africanus* 345-346.
 Lesser Antilles, *Anolis aeneus* (several islands) 264-271.
 Louisiana, *Fundulus (chrysotus, grandis)* 90-92; *Anolis carolinensis* 465-471, 623-624.
 Marshall Islands, *Carcharhinus menisorrhah* 76-83; *Amphiprion* spp., *Chaetodon unimaculatus* 148; *Pomacentrus exilis* 565-568.
 Massachusetts, *Notophthalmus viridescens* 641-647.
 Mexico, *Micrurus diastema* 3; *Poecilia mexicana* (m. mexicana, m. limantouri) 225-236; *Gambusia (atrora, aurata, hurtadoi, krumholzi, marshi, regani, senilis, vittata, panuco)* 250-263; *Poecilia (formosa, mexicana, and hybrids)* 350-352; *Ctenosaura pectinata* 431-434; *Chirostoma* (18 species) 533-554; *Gobiosoma (illecebrosus, prochilos)* 609-610; *Poecilia (formosa, mexicana)* 810-813.
 Minnesota, Forty-two cyprinid species 45-52.
 Mississippi, *Hyla (avivoca, chrysoscelis, cinerea, femoralis, gratiosa, squirella, and various hybrid crosses)* 411-416.
 Missouri, *Cryptobranchus alleganiensis* 331-337; *Eurycea (longicauda melanopleura, lucifuga)* 340-341; *Carphophis vermis* 472-476; *Cryptobranchus alleganiensis bishopi* 805-807.
 Nebraska, *Gerrhonotus* sp., ?*Sceloporus*, *Phrynosoma* ?*douglassi*, ?*Leiocephalus* (fossils) 698-703.
 Nevada, *Cyprinodon nevadensis pectoralis* 137-140; *Acrocheilus alutaceus* × *Richardsonius balteatus* 395-410; *Cyprinodon diabolis* 610-613; *Crotaphytus wislizenii* 658-661; *Gasterosteus (apodus, doryssus, aculeatus aculeatus, a. microcephalus, a. williamsoni)* 721-734; *Lepidomeda (albiwallis, altivelis, mollispinis)*, *Plagopterus argentissimus* 776-782.
 New Mexico, *Bufo cognatus* 103-114; *Meda fulgida* 776-782.
 New York, *Sternotherus odoratus* 305-314.
 New Zealand, *Gobiomorphus gobioides* 169-170; *Leiopelma hochstetteri* 173-175.
 North Carolina, *Ambloplites rupestris* (crossed with *Pomoxis nigromaculatus*, *Enneacanthus obesus*, *Lepomis gulosus*, *Lepomis macrochirus*, *Micropterus salmoides*) 428-430; *Lampropteltis getulus sticticeps* 497-503.
 Ohio, *Etheostoma (blennioides, nigrum, camurum, caeruleum, flabellare)* 163-165; *Umbra* sp. (fossil) 587-588.
 Oklahoma, *Crotalus atrox* 26-31; *Cyprinodon rubrofluvialis* 68-75; *Lepisosteus oculatus* 119-123; *Elaphe buisi* 574-579; *Amia calva* 653-657.
 Oregon, *Lamptetra minima* 568-573; *Xeneretmus leiops* (off Columbia River mouth) 814-817.
 Pacific Ocean (eastern), *Oligoplites refulgens* (Mexico to Ecuador) 151-154; *Oligoplites saurus inornatus* (Mexico, Costa Rica) 151-154; *Psenes (pellucidus, sio)* (Mexico) 167-169; *Bertella idiomorpha* (Mexico, Gulf of California) 193-199; *Xeneretmus leiops* (off Oregon) 814-817.
 Pacific Ocean (western), *Bertella idiomorpha* (off Japan) 193-199; *Mirolabrichthys (tuka tuka, tuka pascualus, dispar)* (various places from Okinawa to Indonesia) 207-208.
 Paraguay (see South America)
 Peru (see South America)
 Rapa (see French Polynesia)
 Seychelles, *Dinematicthys ilucoeteoides* 32-40; *Mirolabrichthys evansi* 207.
 South America (various countries), *Hyla calcarata* (Brazil, Colombia, Ecuador, Guyana, Peru) 515-532; *Hyla fasciata* (Bolivia, Ecuador, Peru) 515-532; *Hyla geographica* (Bolivia, Brazil, Colombia, Ecuador, Guyana, Peru, Surinam, Trinidad and Tobago, Venezuela) 515-532; *Tupinambis teguixin* (Argentina, Bolivia, Brazil, Colombia, French Guiana, Guyana, Paraguay, Surinam, Trinidad, Venezuela) 740-745; *Tupinambis rufescens* (Argentina, Bolivia, Brazil, Paraguay, Uruguay) 740-745.
 South Dakota, *Bufo cognatus* 322-329; *Vorhisia vulpes* (fossil) 692-697.
 Surinam (also see South America), *Lepidocheilus olivacea* 305-314.

- Tennessee, *Lampetra aepyptera* 135; *Eumeces fasciatus* 178-180.
- Texas, *Scaphiopus (bombifrons, hammondi)* 60-67; *Cyprinodon rubrofluvialis* 68-75; *Lepisosteus oculatus* 119-123; *Opsopoeodus emiliae* 161-163; *Gambusia amistadensis* 210-221; *Gambusia (heterochir, georgei, geiseri, amistadensis, nobilis)* 250-263; *Bufo (cognatus, speciosus)* 322-329; *Aphredoderus sayanus* 377-379; *Cnemidophorus tigris* 446-451; *Bufo valliceps* 490-496; *Cyprinodon elegans* × *Cyprinodon variegatus* 682-691.
- Thailand, *Boiga ocellata* 580-586.
- Tobago (see South America)
- Trinidad (see South America)
- Uruguay (see South America)
- Utah, *Sceloporus graciosus* 284-295; *Rhinichthys osculus* × *Richardsonius balteatus*, *Rhinichthys osculus* × *Rhinichthys cataractae* 395-410; *Lepidomeda mollispinis*, *Plagiopterus argentissimus* 776-782.
- Venezuela (also see South America), *Rivulus stellifer* 783-786.
- Wisconsin, *Esox lucius*, *Perca flavescens*, *Lepomis macrochirus* 124-133; *Lepomis cyanellus* 369-373; *Rana pipiens pipiens* 624-627.
- Wyoming, *Gerrhonotus* sp., ?*Sceloporus*, *Phrynosoma* ?*douglasi*, ?*Leiocephalus* 698-703.
- GEOGRAPHIC RANGES**, *Micrurus diastema* 1-4; *Prionobutis koilomatodon* 141; *Bullisichthys caribbaeus* 149-151; *Psenes (pel-lucidus, sio)* 167-169; *Bertella idiomorpha* 197; *Anthias conspicuus* 200; *Gambusia amistadensis* 216-218; *Eleutherodactylus elassodiscus* 223-224; *Poecilia mexicana* (m. mexicana, m. limantouri) 231-234; *Raja (erinacea, ocellata)* 238; *Daector (quadri-zonatus, schmitti)* 355-356; *Diaphus taaningi* 362-363; *Cyprinodon milleri* 366; *Oligocottus maculosus* 373; *Lampropeltis getulus sticticeps* 498-499; *Hyla (calcarata, fasciata, geographica)* 523, 525-527, 530-532; *Chirostoma* (18 species) 533-554; *Pristipomoides squamimaxillaris* 557; *Triathalassothia glowerensis* 560; *Pomacentrus exilis* 565-566; *Lampetra minima* 570; *Boiga ocellata* 581-582, 586; *Tupinambis (teguixin, rufescens)* 741-744; *Sceloporus woodi* 746-748; *Rivulus stellifer* 783; *Liparis inquilinus* 790; *Parrella* (five species) 818.
- GLANDS**, *Chelonia mydas*, *Sternotherus odoratus* (axillary and inguinal glands) 305-314; *Rasbora daniconius* (seasonal changes in pituitary gland) 348-350; *Sceloporus occidentalis* (effect of parietectomy on thyroid and gonads) 662-668; *Sceloporus occidentalis* (thyroid activity) 662-668.
- GRAFTING**, *Hemidactylus garnotii* (heterograft rejection) 339-340.
- GROWTH**, *Python molurus* (effect of temperature) 171-173; *Sceloporus graciosus* 287-288; *Iguana iguana* (embryonic growth) 296-304; *Notophthalmus viridescens* 641-646.
- HERMAPHRODITISM**, *Omosudis lowei*, *Alepisaurus ferox* 41-44; *Bullisichthys caribbaeus* 149-151.
- HISTOLOGY**, *Omosudis lowei*, *Alepisaurus ferox* (reproductive organs) 41-44; *Bullisichthys caribbaeus* (reproductive structures) 149-151; *Chelonia mydas*, *Sternotherus odoratus* (axillary and inguinal glands) 305-314; *Lepomis cyanellus* (effects of temperature and photoperiod on gonads) 369-373; *Carphophis vermis* (testis and epididymis) 472-476; *Sceloporus occidentalis* (epithelial cell height as a measure of thyroid activity) 668-671; *Brachydanio rerio* (development of Mauthner neurons in embryos and larvae) 673-681.
- HOME RANGES** (see **TERRITORIALITY**)
- HORMONES**, *Rasbora daniconius* (seasonal changes in pituitary) 348-350; *Crotaphytus wislizenii* (effects on color change) 658-661; *Sceloporus occidentalis* (effect of parietectomy on thyroid gland and gonads) 662-668; *Sceloporus occidentalis* (thyroid activity and effect on epithelial cell height) 668-672.
- HYBRIDIZATION**, *Acrocheilus alutaceus* × *Richardsonius balteatus* 395-400; *Ambloplites rupestris* × *Enneacanthus obesus*, *Ambloplites rupestris* × *Lepomis* (*Chaenobryttus*) *gulosus*, *Ambloplites rupestris* × *Lepomis macrochirus*, *Ambloplites rupestris* × *Micropterus salmoides*, *Ambloplites rupestris* × *Pomoxis nigromaculatus* 428-430; *Bufo terrestris* × *Bufo woodhousii fowleri*, *Bufo terrestris americanus* × *Bufo woodhousii woodhousii* 61; *Campostoma anomalum* × *Notropis cornutus* 54; *Chrosomus eos* × *Notropis cornutus*, *Chrosomus erythrogaster* × *Notropis cornutus*, *Chrosomus oreas* × *Notropis cornutus* 54-59; *Clinostomus elongatus* × *Notropis cornutus*, *Clinostomus funduloides* × *Notropis cornutus* 54; *Cyprinodon atrox* × *Cyprinodon elegans*, *Cyprinodon elegans* × *Cyprinodon macularius*, *Cyprinodon elegans* × *Cyprinodon nevadensis*, *Cyprinodon elegans* × *Cyprinodon radiosus* 688; *Cyprinodon elegans* × *Cyprinodon variegatus* 682-691;

Cyprinodon macularius × *Cyprinodon variegatus*, *Cyprinodon nevadensis* × *Cyprinodon variegatus*, *Cyprinodon rubrofluvialis* × *Cyprinodon variegatus*, *Cyprinodon salinus* × *Cyprinodon variegatus* 688; *Gila orcutti* × *Hesperoleucus symmetricus* 55, 58, 417-427; *Hyla avivoca* × *Hyla chrysoscelis*, *Hyla avivoca* × *Hyla cinerea*, *Hyla chrysoscelis* × *Hyla cinerea*, *Hyla chrysoscelis* × *Hyla femoralis*, *Hyla chrysoscelis* × *Hyla gratiosa*, *Hyla chrysoscelis* × *Hyla squirella*, *Hyla cinerea* × *Hyla femoralis*, *Hyla cinerea* × *Hyla squirella*, *Hyla femoralis* × *Hyla squirella* 411-416; *Mylocheilus caurinum* × *Ptychocheilus oregonensis* 399-400; *Oncorhynchus gorbuscha* × *Oncorhynchus keta* 58; *Poecilia formosa* × *Poecilia mexicana* 350-352, 813; *Rhinichthys cataractae* × *Rhinichthys osculus*, *Rhinichthys osculus* × *Richardsonius balteatus* 395, 401-406; *Scaphiopus bombifrons* × *Scaphiopus hammondi* 60-67; *Sceloporus undulatus* × *Sceloporus woodi* 756; *Xiphophorus helleri* × *Xiphophorus maculatus* 159-161.

ILLUSTRATIONS (whole animal), *Micrurus diastema* 7; *Paralichthys californicus* (ambicolored) 23; *Hypospsetta guttulata* (ambicolored) 24; *Dinemathichthys ilucoeteoides* (embryos) 35-37; *Notropis cornutus*, *Chrosomus erythrogaster* 55; *Carcharhinus menisora* 79, 81; *Opistognathus aurifrons* 85; *Petromyzon marinus* 136; *Prionobutis koilomatodon* 142; *Megaprotodon strigangulus* 146; *Chaetodon trifasciatus* 147; *Pseudopleuronectes americanus* 155-156; *Psenes sio*, *Psenes pellucidus* 168; *Bertella idiomorpha* 195; *Anthias conspicuus* 204; *Anthias bouleengeri* 206; *Gambusia amistadensis* 214; *Eleutherodactylus elassodiscus* 224; *Poecilia mexicana limantouri* 229; *Poecilia mexicana mexicana* 233; *Raja* (erinacea and ocellata) 244-247; *Daector quadrizonatus* 356; *Galaxias bongbong* 364; *Oligocottus maculosus* (larvae) 374; *Acrocheilus alutaceus* 396; *Richardsonius balteatus* 396; *Acrocheilus alutaceus* × *Richardsonius balteatus* 396; *Rhinichthys osculus* × *Richardsonius balteatus*, *Rhinichthys osculus*, *Rhinichthys cataractae*, *Rhinichthys cataractae* × *Rhinichthys osculus* 401; *Hyla femoralis* × *Hyla squirella* (tadpole) 413; *Hyla geographica* 521; *Hyla calcarata* 524; *Hyla fasciata* 524; *Pristipomoides squamimaxillaris* 558; *Triathalassothia gloverensis*

561; *Pomacentrus exilis* 566; *Lampetra minima* 569; *Salvelinus fontinalis*, *Salmo trutta*, *Gila bicolor* 569; *Boiga ocellata* 581; *Eupomacentrus pictus* 591; *Eupomacentrus partitus* 592; *Iguana iguana* 621; *Cyprinodon elegans*, *Cyprinodon variegatus*, *Cyprinodon elegans* × *Cyprinodon variegatus* 690; *Merriamella doryssa* (= *Gasterosteus doryssus*) 724; *Gasterosteus williamsoni leptosomus* 724; *Gasterosteus apodus* 724; *Ilisha (megaloptera, melastoma)* 736; *Rivulus stellifer* 784; *Liparis inquilinus* 788-789; *Melamphaes ebelingi* 795; *Echis carinatus* 802; *Boiga trigonatum melanocephala* 802; *Bathyagonus nigripinnis* 816; *Brachydanio rerio* 823.

IMMUNE RESPONSES, *Elaphe* (obsoleta, vulpina), *Coluber constrictor* 504-515.

INTERGRADATION, *Poecilia mexicana*: *mexicana* × *limantouri* 228, 234-236; *Lampropeltis getulus*: *getulus* × *sticticeps* 498, 501-502.

INTRODUCTIONS, *Rana catesbeiana* (San Joaquin Valley, California) 18-22; *Prionobutis koilomatodon* (mode of introduction) 141-144; *Gambusia affinis*, *Lepomis cyanellus*, *Micropterus salmoides* (Cuyama River system, California) 420-427; eleven species of fishes (central California) 478-479.

ISOLATING MECHANISMS, *Scaphiopus bombifrons* and *Scaphiopus hammondi* (mating calls) 60-67.

JAWS, *Latimeria chalumnae* (functional anatomy) 156-158.

KARYOTYPES, *Notropis cornutus*, *Chrosomus erythrogaster*, and hybrids 57-59; *Opsopeodus emiliae* 161-163; *Etheostoma* (blennioides, nigrum, camurum, caeruleum, flabellare) 163-165; *Meda fulgida*, *Lepidomeda* (albivallis, mollispinis, vittata), *Plagopterus argentissimus* 776-782.

KEYS (taxonomic), *Raja* (erinacea and ocellata only) (immatures and adults) 248-249; genus *Tupinambis* (teguixin and rufescens) 745; genus *Parrella* (lucretiae, fusca, ginsburgi, macropteryx, maxillaris) 819.

LARVAE, *Megalops atlanticus* 140-141.

LIGHT, *Amia calva* (effect on rate of air breathing) 653-657.

LIPID CYCLES (see **PHYSIOLOGY**)

MERISTICS, *Micrurus diastema* 3-15; pharyngeal-tooth variation in 42 Minnesota cyp-

rinid species (genera *Cyprinus*, *Carassius*, *Pimephales* [3 spp.], *Hybognathus* [2 spp.], *Camptostoma*, *Dionda*, *Phoxinus* [3 spp.], *Clinostomus*, *Semotilus* [2 spp.], *Rhinichthys* [2 spp.], *Couesius*, *Nocomis*, *Hybopsis* [3 spp.], *Phenacobius*, *Notemigonus*, *Opsopoeodus*, *Notropis* [17 spp.]); *Notropis cornutus*, *Chrosomus erythrogaster*, and hybrids 55-57; *Cyprinodon nevadensis pectoralis* 137-139; *Prionobutis koilomatodon* 141; *Heterodontus bonae-spei* (= *Heterodontus portusjacksoni*) (teeth only) 165-167; *Psenes* (sio, *pellucidus*) 168-169; *Bertella idiomorpha* 195-196; *Anthias* (*boulengeri*, *conspicuus*, *squamipinnis*), *Mirolabrichthys* (*evansi*, *tuka tuka*, *tuka pascalus*, *dispar*) 200-202, 205-207; *Gambusia* (*hurtadoi*, *alvarezi*, *gaigei*, *amistadensis*, *senilis*, *nobilis*) 215; *Poecilia mexicana limantouri* 227-228, 234; *Poecilia mexicana mexicana* 232, 234; *Raja* (*erinacea*, *ocellata*) 245; *Poecilia formosa* (in naturally occurring triploid individuals) 351; *Daector quadrizonatus* 355-356; *Engyophrys senta* 357-358; *Diaphus taaningi* 363; *Oligocottus maculosus* (larvae) 375; *Acrocheilus alutaceus* × *Richardsonius balteatus*, *Rhinichthys osculus* × *Richardsonius balteatus*, *Rhinichthys osculus* × *Rhinichthys cataractae* 397-409; *Gila orcutti* × *Hesperoleucus symmetricus* 425-426; *Lampropeltis* (*getulus sticticeps*, *getulus floridana*) 500-503; *Pristipomoides squamimaxillaris* 557-558; *Triathalassothia* (*gloverensis*, *argentinus*), *Amphichthys* (*cryptocentrus*, *hildebrandi*) 560-564; *Pomacentrus exilis* 566; *Lampetra* (*minima*, *tridentata*, *lethophaga*) 570-572; *Elaphe* (*buisi*, *guttata*, *obsoleta*, *triaspis*, *subocularis*, *vulpina*) 577; *Boiga* (*ocellata*, *cyanea*, *dendrophila*, *multimaculatus*, *ochracea*, *quincunciata*, *trigonata*) 582, 584-586; *Eupomacentrus* (*partitus*, *pictus*) 593; *Phrynosoma* (*coronatum*, *cornutum*, *douglasi*, *m'calli*, *modestum*, *platyrhinos*, *solare*) (teeth only) 701; *Gasterosteus* (*apodus*, *dorysus*, *aculeatus aculeatus*, *aculeatus microcephalus*, *aculeatus williamsoni*) 722-723; *Ilisha* (*megaloptera*, *melastoma*) 736, 738; *Tupinambis* (*teguixin*, *rufescens*) 741-742; *Sceloporus woodi* 750; *Rivulus stellifer* 784; *Liparis inquilinus* 790; *Melamphaes ebelingi* 796-799; *BathYGONUS* (*nigripinnis*, *pentacanthus*) 816; *Thymallus arcticus* 821-822.

METABOLISM (see PHYSIOLOGY)

MIMICRY, *Boiga trigonatum melanocephala* and *Echis carinatus* 801-802.

MORPHOMETRICS, *Micrurus diastema* 10-13;

Notropis cornutus, *Chrosomus erythrogaster*, and hybrids 55-57; *Cyprinodon nevadensis pectoralis* 137-139; *Prionobutis koilomatodon* 141; *Heterodontus bonae-spei* (= *Heterodontus portusjacksoni*) (teeth only) 165-167; *Psenes* (sio, *pellucidus*) 169; *Eleotris gobioides* (= *Gobiomorphus gobioides*) 170; *Bertella idiomorpha* 196-197; *Anthias* (*boulengeri*, *conspicuus*), *Mirolabrichthys* (*evansi*, *tuka tuka*, *tuka pascalus*, *dispar*) 201-202, 206-208; *Gambusia* (*hurtadoi*, *alvarezi*, *gaigei*, *amistadensis*, *senilis*, *nobilis*) 211-213, 215, 217-218; *Eleutherodactylus elassodiscus* 222-223; *Poecilia mexicana limantouri* 227-228, 235; *Poecilia mexicana mexicana* 232, 235; *Raja* (*erinacea*, *ocellata*) 240-245; *Iguana iguana* 296-304; *Poecilia formosa* (in naturally-occurring triploid individuals) 350-352; *Engyophrys senta* 357-358; *Acrocheilus alutaceus* × *Richardsonius balteatus*, *Rhinichthys osculus* × *Richardsonius balteatus*, *Rhinichthys osculus* × *Rhinichthys cataractae* 397-409; hybrids among six species of *Hyla* (*avivoca*, *cinerea*, *gratiosa*, *chrysoscelis*, *femoralis*, *squirella*) (tadpoles only) 414-415; *Lampropeltis getulus sticticeps* 501; *Hyla* (*geographica*, *calcarata*, *fasciata*) 517-518, 523, 529; *Pristipomoides squamimaxillaris* 557-558; *Triathalassothia gloverensis* 561; *Pomacentrus exilis* 566-567; *Lampetra* (*minima*, *tridentata*, *lethophaga*) 570-572; *Eupomacentrus* (*partitus*, *pictus*) 591; *Mustelus californicus* (embryos) 597; *Vorhisia vulpes* (otoliths) 696; *Ilisha* (*megaloptera*, *melastoma*) 736, 738; *Sceloporus woodi* 750; *Rivulus stellifer* 784; *Liparis inquilinus* 790; *Melamphaes ebelingi* 796-797; *Thymallus arcticus* 821-822.

NESTS (see REPRODUCTION)

NEURONS, *Brachydanio rerio* (development of Mauthner neurons in embryos and larvae) 673-682.

ORIENTATION, *Crotalus atrox* (radiotelemetry) 26-31; *Bufo valliceps* (olfactory) 490-496.

OSTEOLOGY, *Latimeria chalumnae* (function of jaws) 156-158; *Bertella idiomorpha*, *Dolopichthys pullatus* (hyomandibular, opercular, and lower jaw bones) 194; *Raja* (*erinacea*, *ocellata*) (cartilaginous features) 245-246; *Acrocheilus alutaceus*, *Richardsonius balteatus*, and hybrids (cranial bones) 399-400; *Richardsonius balteatus*, *Rhinichthys osculus*, and hybrids (cranial bones) 399-400, 405, 408; *Boiga ocellata* (cranium)

- 584-586; Gasterosteidae (partial osteology of five genera) 721-734; related to phylogeny of cypriniform fishes 761-775.
- OTOLITHS**, *Vorhisia vulpes* (upper Cretaceous) 692-698.
- OVOVIVIPARITY**, *Dinemachthys ilucoeteoides* 32-40.
- OXYGEN CONSUMPTION** (see **PHYSIOLOGY**)
- PARASITISM** (including **PARASITES**), *Ablepharus boutonii africanus* 345-346; *Lampetra minima* (on *Salmo trutta* and *Salvelinus fontinalis*) 569.
- PHOTOPERIOD**, *Lepomis cyanellus* (effects on seasonal regression) 369-371; *Anolis carolinensis* (influence on ovarian cycle) 465-471.
- PHYSIOLOGY**, *Fundulus (chrysotus, grandis)* (daily rhythms of plasma chloride) 90-92; *Scaphiopus hammondi*, *Bufo cognatus*, *Rana (catesbeiana, pipiens)* (oxygen consumption under different conditions) 106-113; *Dactylagnus mundus* (oxygen consumption) 118; *Lepisosteus oculatus* (effect of dissolved oxygen on locomotor reactions) 119-123; *Esox lucius*, *Perca flavescens*, *Lepomis macrochirus* (behavioral response to low oxygen concentrations) 124-132; *Natrix erythrogaster* (serum proteins and nutritional state) 176-177; *Iguana iguana* (embryonic growth) 296-304; *Sceloporus occidentalis* (induction of physiological color change) 341-342; *Ctenosaura pectinata* (digestive efficiency) 431-434; *Scaphiopus (couchii, hammondi)* (energy metabolism) 435-444; *Cnemidophorus tigris* (energetics and lipid cycles) 446-451; *Scaphiopus (couchii, hammondi)* (gas exchange beneath ground) 452-459; *Dormitator latifrons* (positive buoyancy and air breathing) 461-464; *Anolis carolinensis* (influence of temperature and photoperiod on ovarian cycle) 465-471; *Elaphe (obsoleta, vulpina)*, *Coluber constrictor* (immune response) 504-514; *Coleonyx variegatus* (temperature tolerance) 615-617; *Amia calva* (effect of temperature and light on rate of air breathing) 653-657; *Crotaphytus wislizenii* (hormonal induction of color change in females) 658-667; *Sceloporus occidentalis* (epithelial cell height as measure of thyroid activity) 668-671; *Masticophis flagellum* (electrocardiograph) 802-805; *Cryptobranchus alleganiensis* (thermal acclimation and tolerance) 805-807; *Fundulus (chrysotus, confluentus)* (differential responses to drought) 808-809.
- PIGMENTATION** (including **COLOR**), *Micrurus diastema* (color pattern) 3, 7-13; *Paralichthys californicus*, *Hypopsetta guttulata* (anomalies, ambicoloration) 22-25; *Dinemachthys ilucoeteoides* (color) 32; *Chrosomus erythrogaster* × *Notropis cornutus* (pattern and color) 56; *Cyprinodon rubrofluvialtilis* (color pattern) 68-69; *Bullisichthys caribbaeus* (color and color pattern) 149-151; *Gambusia amistadensis* (color) 215; *Poecilia mexicana limantouri* (pigmentation) 230; *Gambusia (heterochir, affinis, georgei, geiseri, regani, gaigei, marshi)* (variation in anal spot in females) 250-263; *Sceloporus occidentalis* (induction of physiological color change by epinephrine) 341-342; *Oligocottus maculosus* (pigmentation in embryos) 373-377; *Acrocheilus alutaceus* × *Richardsonius balteatus*, *Rhinichthys osculus* × *Richardsonius balteatus*, *Rhinichthys osculus* × *Rhinichthys cataractae* 395-410; *Hyla* (tadpoles of 13 F₁ hybrid combinations involving six species) 411-416; *Lampropeltis getulus sticticeps* 500; *Hyla (geographica, calcarata, fasciata)* 517, 521-526, 529; *Pristipomoides squamimaxillaris* (color) 558; *Triathalassothia gloverensis* (color) 561-562; *Pomacentrus exilis* (color) 566-567; *Lampetra minima* 573; *Boiga (ocellata, cyanea, dendrophila, multimaclulata, ochracea, quincunciata, trigonata)* 583-585; *Eupomacentrus (partitus, pictus)* 590-593; *Mustelus californicus* (albinism) 595-597; *Crotaphytus wislizenii* (hormonal induction of color change) 658-661; *Cyprinodon elegans* × *Cyprinodon variegatus* 684-692; *Ilisha (megalopectera, melastoma)* 737-738; *Tupinambis (teguixin, rufescens)* 741, 743; *Sceloporus woodi* (color variation) 754; *Rivulus stellifer* (color) 784-785; *Liparis inquilinus* 789-790; *Melamphaes ebelingi* 798.
- PLASMA CHLORIDE**, *Fundulus (chrysotus, grandis)* (variation in concentrations) 90-92.
- POLYPLOIDY**, *Poecilia formosa* (triploidy in a form of *Poecilia* very similar to *P. formosa*) 810-813.
- PREDATION**, *Anolis carolinensis* (simulated) 623-624.
- RADIOTELEMETRY**, *Crotalus atrox* 26-31.
- RELATIONSHIPS** (see **TAXONOMY**)
- REPRODUCTION**, family *Brotulidae* (viviparity and oviparity) 32-33, 35, 38; *Chrosomus erythrogaster*, *Notropis cornutus* (breeding behavior) 58-59; *Cyprinodon rubrofluvialtilis* (behavior) 70-75; *Sauromalus obesus* 93-102;

Lampetra aepyptera (nest construction) 135;
Megalops atlanticus (spawning time) 140-141; *Python molurus* (eggs, time of laying, and clutch size) 171; *Gambusia amistadensis* 220; *Gambusia* (*affinis*, *georgei*, *geiseri*, *regani*, *gaigei*, *marshi*, *heterochir*) (courtship) 250-263; *Sceloporus jarrovi* (reproductive effort and winter activity) 272-276; *Sceloporus graciosus* (reproductive cycle) 286-287; *Anolis carolinensis* (effect of temperature and photoperiod on ovarian cycle) 465-471; *Carphophis vermis* (reproductive cycle) 472-476; *Cyprinodon diabolis* (reproductive cycle) 610-613; *Iguana iguana* (post-nesting movements) 620-622.

REPRODUCTIVE CYCLES (see REPRODUCTION)

RESPIRATION (also see PHYSIOLOGY), family Dactyloscopidae (genera *Dactylagnus*, *Gillellus*, *Dactyloscopus*, *Myxodagnus*) (function of respiratory pump) 115-119; *Latimeria chalumnae* (kinetics of) 813-814.

SCALES (including SQUAMATION), *Scomberoides lysan*, *Oligoplites* (*refulgens*, *saurus inornatus*) (feeding on) 151-154; *Brachydanio rerio* (squamation chronology) 823-824.

SEX RATIOS, *Chrosomus erythrogaster* × *Notropis cornutus* 57; *Sceloporus graciosus* 284, 289; *Gila orcutti* × *Hesperoleucus symmetricus* 420; *Cyprinodon elegans* × *Cyprinodon variegatus* 682.

SEXUAL DIMORPHISM, *Micrurus diastema* 3; *Anthias conspicuus* 204; *Gambusia amistadensis* 211-216; *Eleutherodactylus elassodiscus* 223; *Poecilia mexicana* (*mexicana*, *limantouri*) 229, 233; *Gambusia* (seven species) 250-263; *Lampetra minima* 573; *Cyprinodon* (*elegans*, *variegatus*, and hybrids) 690; *Rivulus stellifer* 786.

SHARK ATTACKS, *Carcharhinus menisorrhah* (agonistic display preceding attack) 77-78.

SOUND PRODUCTION (including CALLS), *Scaphiopus* (*bombifrons*, *hammondi*) (isolating mechanisms, spectrograms, effect of temperature) 60-67; *Gnathonemus petersi* 134-135; *Dendrobates pumilio* (vocalizations in territorial behavior) 277-283; *Hyla* (*calcarata*, *fasciata*, *geographica*) (calls and audio-spectrograms) 519-521; *Rana pipiens pipiens* (oscillographs of release calls) 624-627.

SPERM, *Eumeces egregius* (vaginal storage) 346-347; *Carphophis vermis* (spermatogenic cycle) 472-477.

SURVIVORSHIP, *Sceloporus graciosus* 284-296.

SYNONYMIES, *Micrurus diastema* 2; *Hetero-*

dontus bonae-spei (= *Heterodontus portus-jacksoni*) 165-167; *Eleotris gobioides* (= *Gobiomorphus gobioides*) 169-170; *Anthias* (nominal species of) 203; *Anthias formosus* (= *Anthias boulengeri* new name) 206; *Poecilia mexicana* (*limantouri*, *mexicana*) (nomenclature) 227, 232; *Caranx armatus* (= *Carangoides ciliaris*) 352-354; *Engyophrys ciliaris* (= *Engyophrys senta*) 357-358; *Hyla calcarata* 522; *Hyla geographica* 526; *Pungitius haynesi* (= *Gasterosteus aculeatus*) 588-590; *Merriamella doryssa* (= *Gasterosteus doryssus*) 723; *Gasterosteus williamsoni leptosomus* (= *Gasterosteus doryssus*) 723; *Tupinambis* (*teguixin*, *rufescens*) 741, 743; *Liparis inquilinus* 788; *Melamphaes ebelingi* 795; *Asterotheca* (= *BathYGONUS*) 815-817; *Parrella spilopteryx* (= *Parrella lucretiae*) 818.

SYSTEMATICS (see TAXONOMY)

TAXONOMY (including RELATIONSHIPS), *Micrurus diastema* (including relationships) 1-16; *Paralichthys californicus*, *Hypsopsetta guttulata* (abnormal coloration) 22-25; forty-two North American cyprinid fish species (pharyngeal tooth variation) 45-52; *Chrosomus erythrogaster* × *Notropis cornutus* 54-59; *Opsopocodus emiliae* (karyotype) 161-163; *Etheostoma* (*blennioides*, *nigrum*, *camurum*, *caeruleum*, *flabellare*) (karyotypes) 163-165; *Heterodontus bonae-spei* (= *Heterodontus portusjacksoni*) (clarification of status based on examination of jaws of holotype) 165-167; *Psenes* (*pellucidus*, *sio*) (including relationships) 167-169; *Gobiomorphus gobioides* (clarification of status based on examination of types of *Eleotris gobioides*) 169-170; *Bertella idiomorpha*, *Dolopichthys* (including relationships) 193-199; *Anthias* spp. (particularly *boulengeri* and *conspicuus*), *Mirolabrichthys* (*dispar*, *evansi*, *tuka tuka*, *tuka pascalis*) 200-210; *Gambusia* (*amistadensis* and six other species) (includes relationships) 210-221; *Eleutherodactylus elassodiscus* (includes relationships) 222-225; *Poecilia mexicana* (northeastern Mexico) (includes relationships) 225-236; *Raja* (*erinacea*, *ocellata*) 238-249; *Prionotus* (14 species), *Bellator* (four species) (swimbladder and associated structures) 315-321; *Caranx armatus* (= *Carangoides ciliaris*) 352-354; *Daector quadrizonatus* (recognition as valid species) 355-357; *Engyophrys senta* (status of nominal species *Engyophrys ciliaris*) 357-358; *Aphredoderus sayanus* (yolk circulation

patterns as a taxonomic tool) 377-379; *Acrocheilus alutaceus* × *Richardsonius balteatus*, *Rhinichthys osculus* × *Richardsonius balteatus*, *Rhinichthys osculus* × *Rhinichthys cataractae* 395-410; *Hyla* (hybrid tadpoles involving six species: *cinerea*, *avivoca*, *squirella*, *femoralis*, *gratiosa*, *chrysoscelis*) 411-416; *Ambloplites rupestris* (artificial hybridization with five other centrarchid species) 428-430; *Lampropeltis getulus* (*sticticeps*, *floridana*) (including relationships) 500-503; *Hyla* (*geographica*, *calcarata*, *fasciata*) (including relationships) 515-532; *Pristipomoides squamimaxillaris* 557-559; *Triathalassothia gloverensis* (including relationships) 560-565; *Pomacentrus exilis* (including relationships) 565-568; *Lampetra minima* (including relationships) 568-574; *Elaphe buisi* (including relationships and phylogeny of other fossil *Elaphe* species) 574-580; *Boiga* (*ocellata*, *cynodon*) (including relationships) 580-586; *Umbra* sp. (including relationships; Pleistocene fossil) 587-588; *Pungitius haynesi* (= *Gasterosteus aculeatus*) (including relationships; Pliocene fossil) 588-590; *Eupomacentrus* (*partitus*, *pictus*) (comparison between the two species) 590-593; *Cyprinodon elegans* × *Cyprinodon variegatus* (morphological comparison and ecological factors related to hybridization) 682-692; *Vorhisia vulpes* (redescription of otoliths; upper Cretaceous fossils) 692-698; *Gerrhonotus* sp., ?*Sceloporus*, *Phrynosoma* ?*douglasi*, ?*Leiocephalus* (including relationships; Pliocene) 698-704; *Spathorhynchus fossorium* (including relationships within family Rhineuridae; middle Eocene) 704-721; *Gasterosteus* (*apodus*, *doryssus*, *aculeatus aculeatus*, *a. williamsoni*, *a. microcephalus*) (including relationships; Pliocene and Recent) 721-735; *Ilisha* (*megaloptera*, *melastoma*) (including relationships) 735-739; *Tupinambis* (*teguixin*, *rufescens*) (including relationships) 740-746; *Sceloporus woodi* (in part; including relationships to *Sceloporus undulatus*) 746-761; Cypriniform fishes (phylogeny and relationships based primarily on feeding structures) 761-776; Plagopterin cyprinid fishes (*Meda fulgida*, *Lepidomeda* [*vitata*, *altivelis*, *albiwallis*, *mollispinis*] *Plagopterus argentissimus*) (chromosomes) 776-782; *Rivulus stellifer* (including relationships) 783-787; *Liparis* (*inquilinus*, *liparis*, *atlanticus*) (including relationships) 787-794; *Melamphaes ebelingi* (including relationships) 794-800; *Poecilia* (*formosa*, *mexicana*)

(cytogenetic studies of naturally occurring triploids) 810-813; *Asterothea* (= *Bathyagnus*) (*nigripinnis*, *pentacantha*) (including relationships) 815-817; *Parrella* (*lucetiae*, *macropteryx*, *fusca*, *maxillaris*, *ginsburgi*) 817-819; *Thymallus arcticus* 819-822.

TECHNIQUES, lizards (means of obtaining blood from eye) 338-339; amphibians (subcutaneous acrylic polymer injections for marking) 340-341; *Sceloporus occidentalis* (induction of color change by epinephrine) 341-342; aquatic embryos (apparatus for development in controlled environments) 809-810.

TEETH, North American Cyprinidae (42 species; pharyngeal teeth) 45-52; *Chrosomus erythrogaster*, *Notropis cornutus*, and hybrids (pharyngeal teeth) 56; *Heterodontus bonae-spei* (= *Heterodontus portusjacksoni*) (form and number) 165-167; *Bartella idiomorpha*, *Dolopichthys pullatus* 195; *Gambusia amistadensis* 216; *Raja* (*erinacea*, *ocellata*) 243; *Hyla* (hybrid tadpoles involving six species) 414; *Gerrhonotus* sp., ?*Sceloporus*, *Phrynosoma* ?*douglasi*, ?*Leiocephalus* (Miocene fossils) 698-704.

TEMPERATURE, *Cyprinodon rubrofluviatilis* (spawns between 13-34° C) 68-76; *Scaphiopus hammondi* (effect on metabolism) 103-115; *Python molurus* (effect on growth and development) 171-173; *Sceloporus jarrovi* (effect of basking on embryonic development) 272-277; *Cyprinodon milleri* (thermal environment) 366-369; *Lepomis cyanellus* (effects of temperature and photoperiod on seasonal regression) 369-373; *Coleonyx variegatus* (preference) 615-617; *Notophthalmus viridescens* (effect on growth) 641-647; *Amia calva* (effect on rate of air breathing) 653-657; *Cryptobranchius alleganiensis bishopi* (thermal acclimation and tolerance) 805-807.

TERRITORIALITY (including **HOME RANGE**), *Sauromalus obesus* (home range) 93-102; *Megaprotodon strigangulus*, *Chaetodon* (*trifasciatus*, *triangulum*, *unimaculatus*), *Hypsypops rubicunda*, *Pomacentrus flavicauda*, *Abudefduf zonatus*, *Amphiprion* spp. (duration of residence on "home" reefs) 145-149; *Anolis aeneus* (home range in females) 265; *Dendrobates pumilio* 277-284; *Galaxias bonghong* (home range) 363-366.

TRAINING, *Podocnemis unifilis* (techniques) 181-183; *Sphenodon punctatus* (t-maze) 617-620.

TYPE LOCALITIES (not including new species), *Micrurus diastema* (and junior synonyms) 2;

Eleotris gobioides (= *Gobiomorphus gobioides*) 169; *Anthias formosus* (= *Anthias bou-lengeri* new name) 206; *Poecilia mexicana* (mexicana, limantouri) 227, 232; *Daector quadrizonatus* 355; *Engyophrys (ciliaris, senta)* 357-358; *Cyprinodon milleri* 366; *Lampropeltis getulus sticticeps* 498, 500; *Hyla (calcarata, geographica, fasciata, and synonyms)* 522-523, 526; *Merriamella doryssa* (= *Gasterosteus doryssus*), *Gasterosteus williamsoni leptosomus* (= *Gasterosteus doryssus*) (both from Truckee Formation [early Pliocene] of Nevada) 723; *Ilisha (megaloptera, melastoma)* 736; *Tupinambis (teguixin, rufescens)* 741, 743; *Gobius lucretiae* (= *Parrella lucretiae*) 818.

TYPES (not including holotypes of new species)

holotypes, *Heterodontus bonae-spei* (= *Heterodontus portusjacksoni*) 165; *Poecilia mexicana limantouri* 227; *Thalassophryne* (= *Daector*) *quadrizonatus* 355; *Engyophrys senta* 357; *Engyophrys ciliaris* (= *Engyophrys senta*) 357; *Lampropeltis getulus sticticeps* 500; *Hyla leptoscelis* (= *Hyla calcarata*) 522; *Hyla fasciata* 523; *Hyla geographica* (lost) 526; *Hyla geographica* var. *semilineata* (lost) 526; *Hylella punctatissima* (= *Hyla geographica*) 526; *Centrotelma cryptomelan* (= *Hyla geographica*) 526; *Cophomantis punctillata* (= *Hyla geographica*) 526; *Pomacentrus pictus* 591; *Merriamella doryssa* (= *Gasterosteus doryssus*) 724; *Gasterosteus williamsoni leptosomus* (= *Gasterosteus doryssus*) 724; *Clupea melastoma* (= *Ilisha melastoma*) 738; *Gobius* (= *Parrella*) *lucretiae* 818; *Parrella spilopteryx* (= *Parrella lucretiae*) 818; *Parrella maxillaris* 818.

lectotypes, *Elaps diastema* (= *Micrurus diastema*) 2; *Eleotris gobioides* (= *Gobiomorphus gobioides*) 169-170; *Poecilia mexicana mexicana* 232; *Hyla appendiculata* (= *Hyla geographica*) 526; *Lacerta teguixin* (= *Tupinambis teguixin*) 741.

neotype, *Platygaster* (= *Ilisha*) *megalopterus* 737.

paratypes, *Bertella idiomorpha* 194-195; *Anthias conspicuus* 200; *Gambusia amistan-densis* 211; *Eleutherodactylus elassodiscus* 222; *Poecilia mexicana limantouri* 227; *Pristipomoides squamimaxillaris* 557; *Triathalassothia gloverensis* 560; *Pomacentrus exilis* 565; *Lampetra minima* 569; *Boiga ocellata* 581; *Rivulus stellifer* 784; *Liparis inquilinus* 788-789; *Melamphaes ebelingi* 795-796.

syntypes, *Eleotris gobioides* (= *Gobiomorphus gobioides*) 169-170; *Poecilia mexicana mexicana* 232; *Citula ciliaria* (= *Carangoides ciliaris*) 354; *Hyla appendiculata* (= *Hyla geographica*) 526; *Lacerta teguixin* (= *Tupinambis teguixin*) 742; *Tejus rufescens* (= *Tupinambis rufescens*) 743.

VARIATION (morphological), *Micrurus diastema* 1-16; North American Cyprinidae (pharyngeal tooth variation in 42 species) 45-52; *Eleutherodactylus elassodiscus* 222-223; *Gambusia* (seven species) (anal spot expression in females) 250-263; *Hyla (calcarata, fasciata, geographica)* 523, 525-526, 529; *Boiga ocellata* 584; *Thymallus arcticus* 821; *Eupomacentrus partitus* (pigment) 592.

VENTILATION RATES, *Esox lucius*, *Perca flavescens*, *Lepomis macrochirus* (response to low oxygen concentrations) 124-132.

WINTERKILL, *Esox lucius*, *Perca flavescens*, *Lepomis macrochirus* 124-132.

ZOOGEOGRAPHY, *Poecilia mexicana* (in north-eastern Mexico) 234-236; *Orthodon microlepidotus* 358-360; *Chirostoma* (18 species and six subspecies, all in Mexico) 533-554; *Sceloporus woodi* 755-758.